

**MAT 401: Undergraduate Seminar**  
*Introduction to Enumerative Geometry*  
**Fall 2018**

**Homework Assignment VI**

**Written Assignment due on Tuesday, 12/4, at 1pm in ESS 181**

**Problem L**

Let  $F = F(X, Y, Z)$  be a homogeneous polynomial of degree 2 which is not a product of linear factors. Thus,

$$C \equiv Z(F) \equiv \{[X, Y, Z] \in \mathbb{P}^2 : F(X, Y, Z) = 0\}$$

is a smooth curve of degree 2 in  $\mathbb{P}^2$ . Show that there are homogeneous polynomials  $P_i = P_i(u, v)$  of degree 2 so that the image of the map

$$f: \mathbb{P}^1 \longrightarrow \mathbb{P}^2, \quad [u, v] \longrightarrow [P_0(u, v), P_1(u, v), P_2(u, v)]$$

is the curve  $C$ .

There are at least two ways of going about this. The homogeneous polynomials  $P_0, P_1, P_2$  are determined by 3 coefficients each; the homogeneous polynomial  $F$  is given by 6 coefficients. The requirement  $f(\mathbb{P}^1) \subset Z(F)$  is equivalent to

$$F(P_0(u, v), P_1(u, v), P_2(u, v)) = 0 \quad \forall u, v.$$

The left-hand side of this equation is a homogeneous polynomial of degree  $2 \cdot 2$  in  $u$  and  $v$ . Collecting the coefficients of the various terms  $u^a v^{4-a}$ , one obtains 5 equations in 9 unknowns. The extra  $9-5$  degrees of freedom correspond to the fact that if  $P_0, P_1, P_2$  work, so do the polynomials  $P_i(au+bv, cu+dv)$  for any fixed  $a, b, c, d \in \mathbb{C}$ . This approach is direct, but would be very messy.

Here is another approach. It is based on the following observation. Let  $M \in \text{GL}_3\mathbb{C}$  be an invertible  $3 \times 3$ -matrix; it determines a bijective linear map  $M: \mathbb{C}^3 \longrightarrow \mathbb{C}^3$  and induces a bijective map

$$\bar{M}: \mathbb{P}^2 \longrightarrow \mathbb{P}^2, \quad [v] \longrightarrow [Mv].$$

If  $F = F(X, Y, Z)$  is homogeneous polynomial of degree 2, then so is  $F \circ M$ . Furthermore,  $F$  does not split into linear factors if and only if  $F \circ M$  does not (you can prove this either directly or using the approach of Chapter 2, #8). If  $Z(F) = f(\mathbb{P}^1)$ , then  $Z(F \circ M) = \{\bar{M}^{-1} \circ f\}(\mathbb{P}^1)$ , and  $\bar{M}^{-1} \circ f$  is given by the polynomials  $M^{-1}(P_0 P_1 P_2)^{tr}$ . Thus, it is sufficient to prove the statement with  $F$  replaced by  $F \circ M$  for some  $M \in \text{GL}_3\mathbb{C}$ , perhaps repeating the replacement process several times.

For example, if  $F(X, Y, Z) = X^2 + Y^2 + Z^2$ , we could take

$$M = \begin{pmatrix} 1 & i & 0 \\ 1 & -i & 0 \\ 0 & 0 & i \end{pmatrix}^{-1}$$

This replaces  $X+iY$  with  $X$ ,  $X-iY$  with  $Y$ , and  $Z$  with  $iZ$ , so that  $F$  is replaced with  $XY-Z^2$ . So it is sufficient to do the following steps.

(a) Find  $f$  as above that works for  $F(X, Y, Z) = XY - Z^2$ .

(b) If  $F$  does not split into linear factors, show that there exists  $M \in GL_n \mathbb{C}$  so that  $F \circ M$  is  $X^2 + Y^2 + Z^2$  or  $XY - Z^2$ .

## Discussion Problems for 12/4

### *Counting plane rational curves*

Please read the attached note, *even if you are not presenting*, and make sure to actively participate in the discussion, with questions or comments.

If you are presenting,

(1) State formula (1), recalling what  $n_d$  is.

(2) Describe how you are going to prove it; this is essentially Sections 1 and 2.

(3) Prove the formula; this is Section 3 plus you need to derive formula (1) from (6). If time permits, use (1) to compute a few of the numbers  $n_d$ . What is the analogue of this for  $\mathbb{P}^3$ ?

*Please draw pictures*, more of them than in the note, and do not just copy the formulas!

Some of this material is related to some of the material in Chapter 3 of the book.

Please prepare your presentation ahead of time so that it fits in 1 hour and 10 minutes.

# Counting Plane Rational Curves: a modern approach

Aleksey Zinger

October 4, 2018

Enumerative geometry of algebraic varieties is a field of mathematics that dates back to the nineteenth century. The general goal of this subject is to determine the number of geometric objects that satisfy pre-specified geometric conditions. The objects are typically (complex) curves in a smooth algebraic manifold. Such curves are usually required to represent the given homology class, to have certain singularities, and to satisfy various contact conditions with respect to a collection of subvarieties. One of the most well-known examples of an enumerative problem is

**Question 1** *If  $d$  is a positive integer, what is the number  $n_d$  of degree  $d$  rational curves that pass through  $3d-1$  points in general position in the complex projective plane  $\mathbb{P}^2$ ?*

Since the number of (complex) lines through any two distinct points is one,  $n_1 = 1$ . A little bit of algebraic geometry and topology gives  $n_2 = 1$  and  $n_3 = 12$ . It is far harder to find that  $n_4 = 620$ , but this number was computed as early as the middle of the nineteenth century; see [5, p378].

The higher-degree numbers  $n_d$  remained unknown until the early 1990s, when a recursive formula for the numbers  $n_d$  was announced in [2] and [4]:

$$n_d = \frac{1}{6(d-1)} \sum_{d_1+d_2=d} \left( d_1 d_2 - 2 \frac{(d_1 - d_2)^2}{3d - 2} \right) \binom{3d-2}{3d_1-1} d_1 d_2 n_{d_1} n_{d_2}. \quad (1)$$

The argument of the latter paper is described below. It can also be used to solve the natural generalization of Question 1 to the higher-dimensional projective spaces; see [4, Section 10].

We will define an invariant that counts holomorphic maps into  $\mathbb{P}^2$ . A priori, the number we describe depends on the cross ratio of the chosen four points on a sphere. However, it turns out that this number is well-defined. We use its independence to express this invariant in terms of the numbers  $n_d$  in two different ways. By comparing the two expressions, we obtain (1).

## 1 The moduli space of four marked points on a sphere

Let  $x_0, x_1, x_2$  and  $x_3$  be the four points in  $\mathbb{P}^2$  given by

$$x_0 = [1, 0, 0], \quad x_1 = [0, 1, 0], \quad x_2 = [0, 0, 1], \quad x_3 = [1, 1, 1].$$

We denote by  $H^0(\mathbb{P}^2; \gamma^{*\otimes 2})$  the space of holomorphic sections of the holomorphic line bundle  $\gamma^{*\otimes 2} \rightarrow \mathbb{P}^2$ , or equivalently of the degree 2 homogeneous polynomials in three variables. Let

$$\begin{aligned} \mathcal{U} &= \{([s], x) \in \mathbb{P}H^0(\mathbb{P}^2; \gamma^{*\otimes 2}) \times \mathbb{P}^2 : s(x_i) = 0 \ \forall i = 0, 1, 2, 3; \ s(x) = 0\} \\ &\approx \{([A, B]; [z_0, z_1, z_2]) \in \mathbb{P}^1 \times \mathbb{P}^2 : (A-B)z_0z_1 - Az_1z_2 + Bz_0z_2 = 0\}. \end{aligned}$$

The space  $\mathcal{U}$  is a compact complex manifold of dimension 2.

Let  $\pi : \mathcal{U} \rightarrow \overline{\mathcal{M}}_{0,4} \equiv \mathbb{P}^1$  denote the projection onto the first component. If  $[A, B] \in \overline{\mathcal{M}}_{0,4}$ , the fiber  $\pi^{-1}([A, B])$  is the conic

$$\mathcal{C}_{A,B} = \{[z_0, z_1, z_2] \in \mathbb{P}^2 : (A-B)z_0z_1 - Az_1z_2 + Bz_0z_2 = 0\}.$$

If  $[A, B] \neq [1, 0], [0, 1], [1, 1]$ ,  $\mathcal{C}_{A,B}$  is a smooth complex curve of genus zero; it is a sphere with four distinct marked points. If  $[A, B] = [1, 0], [0, 1], [1, 1]$ ,  $\mathcal{C}_{A,B}$  is a union of two lines. One of the lines contains two of the four points  $x_0, \dots, x_3$ , and the other line passes through the remaining two points. The two lines intersect in a single point. Figure 1 shows the three singular fibers of the projection map  $\pi : \mathcal{U} \rightarrow \overline{\mathcal{M}}_{0,4}$ . The other fibers are smooth conics. The fibers should be viewed as lying in planes orthogonal to the horizontal line in the figure.

The following remarks concerning the family  $\mathcal{U} \rightarrow \overline{\mathcal{M}}_{0,4}$  are not directly relevant for the purposes of the next two sections and can be omitted. If  $[A, B] \in \overline{\mathcal{M}}_{0,4} - \{[1, 0], [0, 1], [1, 1]\}$ ,  $\mathcal{C}_{A,B}$  is a smooth complex curve of genus zero, i.e. it is a sphere holomorphically embedded in  $\mathbb{P}^2$ . Thus, there exists a one-to-one holomorphic map  $f : \mathbb{P}^1 \rightarrow \mathcal{C}_{A,B}$ . It can be shown directly that if  $[u_i, v_i] = f^{-1}(x_i)$ ,

$$\frac{v_0/u_0 - v_2/u_2}{v_0/u_0 - v_3/u_3} : \frac{v_1/u_1 - v_2/u_2}{v_1/u_1 - v_3/u_3} = \frac{B}{A}.$$

The cross-ratio is the only invariant of four distinct points on  $\mathbb{P}^1$ ; see [1], for example. Thus,

$$\begin{aligned} \mathbb{P}^1 - \{[1, 0], [0, 1], [1, 1]\} &= \mathcal{M}_{0,4} \equiv \{(x_0, x_1, x_2, x_3) \in (\mathbb{P}^1)^4 : x_i \neq x_j \text{ if } i \neq j\} / \sim, \\ \text{where } (x_0, x_1, x_2, x_3) &\sim (\tau(x_0), \tau(x_1), \tau(x_2), \tau(x_3)) \quad \text{if } \tau \in \text{PSL}_2 \equiv \text{Aut}(\mathbb{P}^1). \end{aligned}$$

Furthermore, the restriction of the projection map  $\pi : \mathcal{U}|_{\mathcal{M}_{0,4}} \rightarrow \mathcal{M}_{0,4}$  to each fiber  $\mathcal{C}_{[A,B]}$  is the cross ratio of the points  $x_0, \dots, x_3$  on  $\mathcal{C}_{[A,B]}$ , viewed as an element of  $\mathbb{P}^1 \supset \mathbb{C}$ .

## 2 Counts of holomorphic maps

If  $d$  is an integer and  $\mathcal{C}$  is a complex curve, which may be a wedge of spheres, let

$$\mathcal{H}_d(\mathcal{C}) = \{f \in C^\infty(\mathcal{C}; \mathbb{P}^2) : f \text{ is holomorphic, } \deg f = d\}. \quad (2)$$

We give a more explicit description of the space  $\mathcal{H}_d(\mathcal{C})$  in the relevant cases below.

Suppose  $\ell_0, \ell_1$  and  $p_2, \dots, p_{3d-1}$  are two lines and  $3d-2$  points in general position in  $\mathbb{P}^2$ . If  $\sigma \in \overline{\mathcal{M}}_{0,4}$ , let  $N_d^\sigma(\ell_0, \ell_1, p_2, \dots, p_{3d-1})$  denote the cardinality of the set

$$\{f \in \mathcal{H}_d(\mathcal{C}_\sigma) : f(x_0) \in \ell_0, f(x_1) \in \ell_1, f(x_2) = p_2, f(x_3) = p_3, p_i \in \text{Im } f \ \forall i\}. \quad (3)$$




Figure 1: The Family  $\mathcal{U} \rightarrow \overline{\mathcal{M}}_{0,4}$

Here  $\mathcal{C}_\sigma$  denotes the rational curve with four marked points,  $x_0, x_1, x_2$ , and  $x_3$ , whose cross ratio is  $\sigma$ ; see Section 1. If  $\sigma \neq [1, 0], [0, 1], [1, 1]$ ,  $\mathcal{C}_\sigma$  is a sphere with four, distinct, marked points. In this case, the condition  $f \in \mathcal{H}_d(\mathcal{C}_\sigma)$  means that  $f$  has the form

$$f([u, v]) = [P_0(u, v), P_1(u, v), P_2(u, v)] \quad \forall [u, v] \in \mathbb{P}^1,$$

for some degree  $d$  homogeneous polynomials  $P_0, P_1, P_2$  that have no common factor. If  $\sigma = [1, 0], [0, 1], [1, 1]$ ,  $\mathcal{C}_\sigma$  is a wedge of two spheres,  $\mathcal{C}_{\sigma,1}$  and  $\mathcal{C}_{\sigma,2}$ , with two marked points each. In this case, the first condition in (2) means that  $f$  is continuous and  $f|_{\mathcal{C}_{\sigma,1}}$  and  $f|_{\mathcal{C}_{\sigma,2}}$  are holomorphic. The second condition in (2) means that  $d = d_1 + d_2$  if the degrees of  $f|_{\mathcal{C}_{\sigma,1}}$  and  $f|_{\mathcal{C}_{\sigma,2}}$  are  $d_1$  and  $d_2$ , respectively.

The requirement that the two lines,  $\ell_0$  and  $\ell_1$ , and the  $3d-2$  points,  $p_2, \dots, p_{3d-1}$ , are in general position means that they lie in a dense open subset  $\mathcal{U}_\sigma$  of the space of all possible tuples  $(\ell_0, \ell_1, p_2, \dots, p_{3d-1})$ :

$$\mathfrak{X} \equiv \text{Gr}_2\mathbb{C}^3 \times \text{Gr}_2\mathbb{C}^3 \times (\mathbb{P}^2)^{3d-2}.$$

Here  $\text{Gr}_2\mathbb{C}^3$  denotes the Grassmanian manifold of two-planes through the origin in  $\mathbb{C}^3$ , or equivalently of lines in  $\mathbb{P}^2$ . The dense open subset  $\mathcal{U}_\sigma$  of  $\mathfrak{X}$  consists of tuples  $(\ell_0, \ell_1, p_2, \dots, p_{3d-1})$  that satisfy a number of geometric conditions. In particular,  $\ell_0 \neq \ell_1$ , none of the points  $p_2, \dots, p_{3d-1}$  lies on either  $\ell_0$  or  $\ell_1$ , the  $3d-1$  points  $\ell_0 \cap \ell_1, p_2, \dots, p_{3d-1}$  are distinct, no three of them lie on the same line, and so on. In addition, we need to impose certain cross-ratio conditions on the rational curves that pass through  $\ell_0, \ell_1, p_2, p_3$ , and a subset of the remaining  $3d-4$  points. These conditions can be stated more formally. Define

$$\text{ev}_\sigma: \mathcal{H}_d(\mathcal{C}_\sigma) \times (\mathcal{C}_\sigma)^{3d-4} \rightarrow (\mathbb{P}^2)^{3d} \quad \text{by} \quad \text{ev}_\sigma(f; x_4, \dots, x_{3d-1}) = (f(x_0), f(x_1), \dots, f(x_{3d-1})).$$

The space  $\mathcal{H}_d(\mathcal{C}_\sigma)$  is a dense open subset of  $\mathbb{P}^{3d+2}$  and the evaluation map  $\text{ev}_\sigma$  is holomorphic. There is a natural compactification  $\overline{\mathfrak{M}}_\sigma(\mathbb{P}^2, d)$  of  $\mathcal{H}_d(\mathcal{C}_\sigma)$ , which consists spaces of holomorphic maps from various wedges of spheres into  $\mathbb{P}^2$ . The complex dimension of each such boundary stratum is less than that of  $\mathcal{H}_d(\mathcal{C}_\sigma)$ . The evaluation map  $\text{ev}_\sigma$  admits a continuous extension over  $\partial\overline{\mathfrak{M}}_\sigma(\mathbb{P}^2, d)$ , whose restriction to each stratum is holomorphic. The elements  $(\ell_0, \ell_1, p_2, \dots, p_{3d-1})$  of the subspace  $\mathcal{U}_\sigma$  of  $\mathfrak{X}$  are characterized by the condition that the restriction of the evaluation map to each stratum of  $\overline{\mathfrak{M}}_\sigma(\mathbb{P}^2, d)$  is transversal to the submanifold

$$\ell_0 \times \ell_1 \times p_2 \times \dots \times p_{3d-1} \subset (\mathbb{P}^2)^{3d}.$$

This condition implies that

$$\text{ev}_\sigma^{-1}(\ell_0 \times \ell_1 \times p_2 \times \dots \times p_{3d-1}) \cap \partial \overline{\mathfrak{M}}_\sigma(\mathbb{P}^2, d) = \emptyset$$

and the set in (3) is a finite subset of  $\mathcal{H}_d(\mathcal{C}_\sigma)$ .

The set  $\mathcal{U}_\sigma$  of “general” tuples  $(\ell_0, \ell_1, p_2, \dots, p_{3d-1})$  is path-connected. Indeed, it is the complement of a finite number of proper complex submanifolds in  $\mathfrak{X}$ . It follows that the number in (3) is independent of the choice of two lines and  $3d-2$  points in general position in  $\mathbb{P}^2$ . We thus may simply denote it by  $N_d^\sigma$ . If  $\sigma \neq [1, 0], [0, 1], [1, 1]$ ,  $\mathcal{C}_\sigma$  is a sphere with four distinct points. In such a case, it is fairly easy to show that the number  $N_d^\sigma$  does not change under small variations of  $\sigma$ , or equivalently of the four points on the sphere. Thus,  $N_d^\sigma$  is independent of

$$\sigma \in \mathcal{M}_{0,4} = \mathbb{P}^1 - \{[1, 0], [0, 1], [1, 1]\} = \overline{\mathcal{M}}_{0,4} - \{[1, 0], [0, 1], [1, 1]\}.$$

It is far harder to prove

**Proposition 2** *The function  $\sigma \rightarrow N_d^\sigma$  is constant on  $\overline{\mathcal{M}}_{0,4}$ .*

This proposition is a special case of the gluing theorems first proved in [3] and [4].

### 3 Holomorphic maps vs. complex curves

In this subsection, we express the numbers  $N_d^{[0,1]}$  and  $N_d^{[1,1]}$  of Subsections 2 in terms of the numbers  $n_{d'}$ , with  $d' \leq d$ , of Question 1. By Proposition 2,  $N_d^{[0,1]} = N_d^{[1,1]}$ . We obtain a recursion for the numbers of Question 1 by comparing the expressions for  $N_d^{[0,1]}$  and  $N_d^{[1,1]}$ .

Let  $\mathcal{C}_1$  denote the component of  $\mathcal{C}_{[0,1]}$  containing the marked points  $x_0$  and  $x_3$ ; see Figure 1. We denote by  $\mathcal{C}_2$  the other component of  $\mathcal{C}_{[0,1]}$ . By definition,

$$N_d^{[0,1]} = \sum_{d_1+d_2=d} N_{d_1, d_2}^{[0,1]} \quad \text{where}$$

$$N_{d_1, d_2}^{[0,1]} = |\{f \in \mathcal{H}_d(\mathcal{C}_{[0,1]}; \mathbb{P}^2) : \deg f|_{\mathcal{C}_1} = d_1, \deg f|_{\mathcal{C}_2} = d_2; p_i \in \text{Im } f \forall i; f(x_0) \in \ell_0, f(x_1) \in \ell_1, f(x_2) = p_2, f(x_3) = p_3\}|.$$

Since the group  $PSL_2$  of holomorphic automorphisms acts transitively on triples of distinct points on the sphere,

$$N_{d_1, d_2}^{[0,1]} = |\{(f_1, f_2) \in \mathcal{H}_{d_1}(S^2) \times \mathcal{H}_{d_2}(S^2) : f_1(\infty) = f_2(\infty), p_i \in f_1(S^2) \cup f_2(S^2) \forall i; f_1(0) \in \ell_0, f_1(1) = p_3, f_2(0) \in \ell_1, f_2(1) = p_2\}|.$$

Since the maps  $f_1$  and  $f_2$  above are holomorphic,  $d_1, d_2 \geq 0$  if  $N_{d_1, d_2}^{[0,1]} \neq 0$ . Since every degree 0 holomorphic map is constant and  $p_3 \notin \ell_0$ ,  $N_{0, d}^{[0,1]} = 0$ . Similarly,  $N_{d, 0}^{[0,1]} = 0$ . Thus, we assume that  $d_1, d_2 > 0$ . Since the points  $p_3, \dots, p_{3d-1}$  are in general position,  $f_1(S^2)$  contains at most  $3d_1 - 2$  of

the points  $p_4, \dots, p_{3d-1}$ . Similarly, the curve  $f_2(S^2)$  passes through at most  $3d_2 - 2$  of the points  $p_4, \dots, p_{3d-1}$ . Thus, if  $I = \{4, \dots, 3d-1\}$ ,

$$N_{d_1, d_2}^{[0,1]} = \sum_{I=I_1 \sqcup I_2, |I_1|=3d_1-2} N_{d_1, d_2}^{[0,1]}(I_1, I_2),$$

where  $N_{d_1, d_2}^{[0,1]}(I_1, I_2)$  is the cardinality of the set

$$\begin{aligned} \mathcal{S}_{d_1, d_2}^{[0,1]}(I_1, I_2) = \{ & (f_1, f_2) \in \mathcal{H}_{d_1}(S^2) \times \mathcal{H}_{d_2}(S^2) : p_i \in f_1(S^2) \ \forall i \in I_1, \ p_i \in f_2(S^2) \ \forall i \in I_2; \\ & f_1(\infty) = f_2(\infty), \ f_1(0) \in \ell_0, \ f_1(1) = p_3, \ f_2(0) \in \ell_1, \ f_2(1) = p_2\}. \end{aligned}$$

If  $(f_1, f_2) \in \mathcal{S}_{d_1, d_2}^{[0,1]}(I_1, I_2)$ ,  $f_1(S^2)$  is one of the  $n_{d_1}$  curves passing through the points  $\{p_i : i \in \{3\} \sqcup I_1\}$ . Similarly,  $f_2(S^2)$  is one of the  $n_{d_2}$  curves passing through the points  $\{p_i : i \in \{2\} \sqcup I_2\}$ . The point  $f_1(\infty) = f_2(\infty)$  must be one of the points of  $f_1(S^2) \cap f_2(S^2)$ ; by Bezout's theorem there are  $d_1 d_2$  such points. Finally,  $f_1(0)$  must be one of the  $d_1$  points of  $f_1(S^2) \cap \ell_0$ , while  $f_2(0)$  must be one of the  $d_2$  points of  $f_2(S^2) \cap \ell_1$ . Thus, we conclude that

$$\begin{aligned} N_d^{[0,1]} &= \sum_{d_1+d_2=d} N_{d_1, d_2}^{[0,1]} = \sum_{d_1+d_2=d} \sum_{I=I_1 \sqcup I_2, |I_1|=3d_1-2} N_{d_1, d_2}^{[0,1]}(I_1, I_2) \\ &= \sum_{d_1+d_2=d} \sum_{I_1 \subset I, |I_1|=3d_1-2} (d_1 d_2) (d_1 n_{d_1}) (d_2 n_{d_2}) \\ &= \sum_{d_1+d_2=d} \binom{3d-4}{3d_1-2} d_1^2 d_2^2 n_{d_1} n_{d_2}; \end{aligned} \tag{4}$$

where  $I = \{4, \dots, 3d-1\}$ .

We compute the number  $N_d^{[1,1]}$  similarly. We denote by  $\mathcal{C}_1$  the component of  $\mathcal{C}_{[1,1]}$  containing the points  $x_0$  and  $x_1$  and by  $\mathcal{C}_2$  the other component of  $\mathcal{C}_{[1,1]}$ . By definition,

$$\begin{aligned} N_d^{[1,1]} &= \sum_{d_1+d_2=d} N_{d_1, d_2}^{[1,1]}, \quad \text{where} \\ N_{d_1, d_2}^{[1,1]} &= \left| \left\{ (f_1, f_2) \in \mathcal{H}_{d_1}(S^2) \times \mathcal{H}_{d_2}(S^2) : f_1(\infty) = f_2(\infty), \ p_i \in f_1(S^2) \cup f_2(S^2) \ \forall i; \right. \right. \\ & \quad \left. \left. f_1(0) \in \ell_0, \ f_1(1) \in \ell_1, \ f_2(0) = p_2, \ f_2(1) = p_3 \right\} \right|. \end{aligned}$$

Since every degree-zero holomorphic map is constant,  $N_{d,0}^{[1,1]} = 0$  as before. However,

$$\begin{aligned} N_{0,d}^{[1,1]} &= \left| \left\{ f_2 \in \mathcal{H}_d(S^2) : f_2(\infty) \in \ell_0 \cap \ell_1, \ f_2(0) = p_2, \ f_2(1) = p_3; \right. \right. \\ & \quad \left. \left. p_i \in f_2(S^2) \ \forall i = 4, \dots, 3d-1 \right\} \right|. \end{aligned}$$

Thus,  $N_{0,d}^{[1,1]} = n_d$ . If  $d_1, d_2 > 0$ ,

$$N_{d_1, d_2}^{[1,1]} = \sum_{I=I_1 \sqcup I_2, |I_1|=3d_1-1} N_{d_1, d_2}^{[1,1]}(I_1, I_2),$$

where  $N_{d_1, d_2}^{[1,1]}(I_1, I_2)$  is the cardinality of the set

$$\mathcal{S}_{d_1, d_2}^{[1,1]}(I_1, I_2) = \{(f_1, f_2) \in \mathcal{H}_{d_1}(S^2) \times \mathcal{H}_{d_2}(S^2) : p_i \in f_1(S^2) \forall i \in I_1, p_i \in f_2(S^2) \forall i \in I_2; \\ f_1(\infty) = f_2(\infty), f_1(0) \in \ell_0, f_1(1) \in \ell_1, f_2(0) = p_2, f_2(1) = p_3\}.$$

Proceeding as in the previous paragraph, we conclude that

$$\begin{aligned} N_d^{[1,1]} &= \sum_{d_1+d_2=d} N_{d_1, d_2}^{[1,1]} = n_d + \sum_{d_1+d_2=d} \sum_{I=I_1 \sqcup I_2, |I_1|=3d_1-1} N_{d_1, d_2}^{[1,1]}(I_1, I_2) \\ &= n_d + \sum_{d_1+d_2=d} \sum_{I_1 \subset I, |I_1|=3d_1-1} (d_1 d_2) (d_1^2 n_{d_1}) (n_{d_2}) \\ &= n_d + \sum_{d_1+d_2=d} \binom{3d-4}{3d_1-1} d_1^3 d_2 n_{d_1} n_{d_2}; \end{aligned} \tag{5}$$

Comparing equations (4) and (5), we obtain

$$n_d = \sum_{d_1+d_2=d} \left( \binom{3d-4}{3d_1-2} d_1 d_2 - \binom{3d-4}{3d_1-1} d_1^2 \right) d_1 d_2 n_{d_1} n_{d_2}. \tag{6}$$

The recursive formula (1) is the symmetrized version of (6).

## References

- [1] L. Ahlfors, *Complex Analysis*, McGraw-Hill, 1979.
- [2] M. Kontsevich and Yu. Manin, *Gromov-Witten Classes, Quantum Cohomology, and Enumerative Geometry*, Comm. Math. Phys. 164 (1994), no. 3, 525–562.
- [3] D. McDuff and D. Salamon, *Introduction to J-Holomorphic Curves*, American Mathematical Society, 1994.
- [4] Y. Ruan and G. Tian, *A Mathematical Theory of Quantum Cohomology*, J. Diff. Geom. 42 (1995), no. 2, 259–367.
- [5] H. Zeuthen, *Almindelige Egenskaber ved Systemer af Plane Kurver*, Kongelige Danske Videnskaberne Selskabs Skrifter, 10 (1873), 285–393. Danish.