

MAT 542: Algebraic Topology, Fall 2016

Suggested Problems for Week 11

You may hand in solutions to at *most* 2 problems every 2 weeks and no later than 2 weeks after the necessary material for them is covered in class.

From Munkres: 47.1, 47.3, 47.5, 47.7, 48.1-3, 49.1-3, 49.8, 66.4-6, 57.1

Problem Q

Let R be a commutative ring with 1 and M be a module over R . A **directed set** is a set S with a partial order \prec such that for any pair of elements $i, j \in S$ there exists $k \in S$ with $i, j \leq k$. A **directed system** of R -modules over a directed set (S, \prec) is an assignment of an R -module M_i for each $i \in S$ and an R -module homomorphism $\rho_{ji}: M_i \rightarrow M_j$ for all $i, j \in S$ with $i \leq j$ such that

$$\rho_{ii} = \text{id}_{M_i} \quad \forall i \in S, \quad \rho_{ki} = \rho_{kj} \circ \rho_{ji} \quad \forall i, j, k \in S, \quad i \leq j \leq k.$$

An **inverse system** of R -modules over a directed set (S, \prec) is an assignment of an R -module M_i for each $i \in S$ and an R -module homomorphism $\varrho_{ij}: M_j \rightarrow M_i$ for all $i, j \in S$ with $i \leq j$ such that

$$\varrho_{ii} = \text{id}_{M_i} \quad \forall i \in S, \quad \varrho_{ik} = \varrho_{ij} \circ \varrho_{jk} \quad \forall i, j, k \in S, \quad i \leq j \leq k.$$

The direct limit of a directed system and the inverse limit of an inverse system are

$$\varinjlim M_i \equiv \left(\bigsqcup_i M_i \right) / \sim, \quad A_i \ni x_i \sim x_j \in A_j \quad \text{if } \exists k \in S \text{ s.t. } i, j \leq k, \quad \rho_{ki}(x_i) = \rho_{kj}(x_j),$$

$$\varprojlim M_i \equiv \left\{ (x_i)_{i \in S} \in \prod_i M_i : x_i = \varrho_{ij}(x_j) \quad \forall i, j \in S, \quad i \prec j \right\},$$

respectively. For example, the collection of compact subsets K of a topological spaces forms a directed set. The associated homology and cohomology groups form a direct system and an inverse system under the homomorphisms

$$\rho_{K'K}: H_*(K; R) \rightarrow H_*(K'; R) \quad \text{and} \quad \varrho_{KK'}: H^*(K'; M) \rightarrow H^*(K; M)$$

induced by the inclusions $K \subset K'$.

- (a) Show that the direct and inverse limits are R -modules.
- (b) Construct isomorphisms

$$\varinjlim H_*(K; R) \rightarrow H_*(X; R), \quad H_*(X; R) \rightarrow \varprojlim H^*(K; R).$$

Problem R

Let R be a commutative ring with 1. Let X, Y be Hausdorff topological spaces and $X \vee Y$ be their wedge product at points that are deformation retracts of neighborhoods in their spaces.

- (a) Construct an isomorphism

$$H^p(X \vee Y; R) \rightarrow H^p(X; R) \oplus H^p(Y; R) \quad \forall p \in \mathbb{Z}^+$$

and show that it respects the cup product.

- (b) Use this to establish the claims of Example 4 and Exercise 5 in Section 49 without any simplicial complexes.