

Instantons, Quotient Singularities,
and the
Geometrization of Four-Manifolds

Claude LeBrun
Stony Brook University

Vienna, Sept 10, 2012



For Mike Eastwood



For Mike Eastwood

who taught me
never to be afraid of
a Dynkin diagram





For Mike Eastwood

who taught me
never to be afraid of
a Dynkin diagram


but who, somehow,
never convinced me to
try bungee jumping.

The Dynkin diagrams of simply laced Lie groups

The Dynkin diagrams of simply laced Lie groups




The Dynkin diagrams of simply laced Lie groups




play an important role in 4-dimensional geometry,

The Dynkin diagrams of simply laced Lie groups



play an important role in 4-dimensional geometry,
particularly in connection with Einstein manifolds.


The Dynkin diagrams of simply laced Lie groups



play an important role in 4-dimensional geometry,
particularly in connection with Einstein manifolds.

This might be dismissed as a combinatorial fluke.

The Dynkin diagrams of simply laced Lie groups



play an important role in 4-dimensional geometry,
particularly in connection with Einstein manifolds.

This might be dismissed as a combinatorial fluke.

But it is directly related to representation theory.

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.


“... the greatest blunder of my life!”

— A. Einstein, to G. Gamow

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.



Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

As punishment ...

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

λ called Einstein constant.

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

λ called Einstein constant.

Has same sign as the *scalar curvature*

$$s = r_j^j = \mathcal{R}^{ij}_{ij}.$$

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$


for some constant $\lambda \in \mathbb{R}$.

λ called Einstein constant.

Has same sign as the *scalar curvature*

$$s = r_j^j = \mathcal{R}^{ij}_{ij}.$$

$$\frac{\text{vol}_g(B_\varepsilon(p))}{c_n \varepsilon^n} = 1 - s \frac{\varepsilon^2}{6(n+2)} + O(\varepsilon^4)$$



Variational Problem:

Variational Problem:

If M smooth compact n -manifold,

Variational Problem:

If M smooth compact n -manifold, $n \geq 3$,

Variational Problem:

If M smooth compact n -manifold, $n \geq 3$,

$$\mathcal{G}_M = \{ \text{smooth metrics } g \text{ on } M \}$$

Variational Problem:

If M smooth compact n -manifold, $n \geq 3$,

$$\mathcal{G}_M = \{ \text{smooth metrics } g \text{ on } M \}$$

then Einstein metrics are critical points

Variational Problem:

If M smooth compact n -manifold, $n \geq 3$,

$$\mathcal{G}_M = \{ \text{smooth metrics } g \text{ on } M \}$$

then Einstein metrics are critical points of the scale-invariant action functional

Variational Problem:

If M smooth compact n -manifold, $n \geq 3$,

$$\mathcal{G}_M = \{ \text{smooth metrics } g \text{ on } M \}$$

then Einstein metrics are critical points of the scale-invariant action functional

$$\begin{aligned} \mathcal{G}_M &\longrightarrow \mathbb{R} \\ g &\longmapsto \int_M |s_g|^{n/2} d\mu_g \end{aligned}$$

Variational Problem:

If M smooth compact n -manifold, $n \geq 3$,

$$\mathcal{G}_M = \{ \text{smooth metrics } g \text{ on } M \}$$

then Einstein metrics are critical points of the scale-invariant action functional

$$\begin{aligned} \mathcal{G}_M &\longrightarrow \mathbb{R} \\ g &\longmapsto \int_M |s_g|^{n/2} d\mu_g \end{aligned}$$

Conversely:

Variational Problem:

If M smooth compact n -manifold, $n \geq 3$,

$$\mathcal{G}_M = \{ \text{smooth metrics } g \text{ on } M \}$$

then Einstein metrics are critical points of the scale-invariant action functional

$$\begin{aligned} \mathcal{G}_M &\longrightarrow \mathbb{R} \\ g &\longmapsto \int_M |s_g|^{n/2} d\mu_g \end{aligned}$$

Conversely:

Weak critical points are Einstein

Variational Problem:

If M smooth compact n -manifold, $n \geq 3$,

$$\mathcal{G}_M = \{ \text{smooth metrics } g \text{ on } M \}$$

then Einstein metrics are critical points of the scale-invariant action functional

$$\begin{aligned} \mathcal{G}_M &\longrightarrow \mathbb{R} \\ g &\longmapsto \int_M |s_g|^{n/2} d\mu_g \end{aligned}$$

Conversely:

Weak critical points are Einstein or scalar-flat ($s \equiv 0$).

Variational Problem:

If M smooth compact n -manifold, $n \geq 3$,

$$\mathcal{G}_M = \{ \text{smooth metrics } g \text{ on } M \}$$

then Einstein metrics are critical points of the scale-invariant action functional

$$\begin{aligned} \mathcal{G}_M &\longrightarrow \mathbb{R} \\ g &\longmapsto \int_M |s_g|^{n/2} d\mu_g \end{aligned}$$

Conversely:

Weak critical points are Einstein or scalar-flat ($s \equiv 0$).

Try to find Einstein metrics by minimizing?

A Differential-Topological Invariant:

A Differential-Topological Invariant:

$$\int_M |s_g|^{n/2} d\mu_g$$

A Differential-Topological Invariant:

$$\inf_g \int_M |s_g|^{n/2} d\mu_g$$

A Differential-Topological Invariant:

$$\mathcal{I}_s(M) = \inf_g \int_M |s_g|^{n/2} d\mu_g$$

A Differential-Topological Invariant:

$$\mathcal{I}_s(M) = \inf_g \int_M |s_g|^{n/2} d\mu_g$$

Theorem. *Let M be a compact simply connected n -manifold,*

A Differential-Topological Invariant:

$$\mathcal{I}_s(M) = \inf_g \int_M |s_g|^{n/2} d\mu_g$$

Theorem. *Let M be a compact simply connected n -manifold, $n \geq 3$.*

A Differential-Topological Invariant:

$$\mathcal{I}_s(M) = \inf_g \int_M |s_g|^{n/2} d\mu_g$$

Theorem. *Let M be a compact simply connected n -manifold, $n \geq 3$. If $n \neq 4$,*

A Differential-Topological Invariant:

$$\mathcal{I}_s(M) = \inf_g \int_M |s_g|^{n/2} d\mu_g$$

Theorem. *Let M be a compact simply connected n -manifold, $n \geq 3$. If $n \neq 4$, $\mathcal{I}_s(M) = 0$.*

A Differential-Topological Invariant:

$$\mathcal{I}_s(M) = \inf_g \int_M |s_g|^{n/2} d\mu_g$$

Theorem. *Let M be a compact simply connected n -manifold, $n \geq 3$. If $n \neq 4$, $\mathcal{I}_s(M) = 0$.*

Theorem. *There exist compact simply connected 4-manifolds M_j*

A Differential-Topological Invariant:

$$\mathcal{I}_s(M) = \inf_g \int_M |s_g|^{n/2} d\mu_g$$

Theorem. *Let M be a compact simply connected n -manifold, $n \geq 3$. If $n \neq 4$, $\mathcal{I}_s(M) = 0$.*

Theorem. *There exist compact simply connected 4-manifolds M_j with $\mathcal{I}_s(M_j) \rightarrow +\infty$.*

A Differential-Topological Invariant:

$$\mathcal{I}_s(M) = \inf_g \int_M |s_g|^{n/2} d\mu_g$$

Theorem. *Let M be a compact simply connected n -manifold, $n \geq 3$. If $n \neq 4$, $\mathcal{I}_s(M) = 0$.*

Theorem. *There exist compact simply connected 4-manifolds M_j with $\mathcal{I}_s(M_j) \rightarrow +\infty$.*

Moreover, can choose M_j

A Differential-Topological Invariant:

$$\mathcal{I}_s(M) = \inf_g \int_M |s_g|^{n/2} d\mu_g$$

Theorem. Let M be a compact simply connected n -manifold, $n \geq 3$. If $n \neq 4$, $\mathcal{I}_s(M) = 0$.

Theorem. There exist compact simply connected 4-manifolds M_j with $\mathcal{I}_s(M_j) \rightarrow +\infty$.

Moreover, can choose M_j such that

$$\mathcal{I}_s(M_j) = \inf_g \int_{M_j} |s_g|^2 d\mu_g$$

A Differential-Topological Invariant:

$$\mathcal{I}_s(M) = \inf_g \int_M |s_g|^{n/2} d\mu_g$$

Theorem. Let M be a compact simply connected n -manifold, $n \geq 3$. If $n \neq 4$, $\mathcal{I}_s(M) = 0$.

Theorem. There exist compact simply connected 4-manifolds M_j with $\mathcal{I}_s(M_j) \rightarrow +\infty$.

Moreover, can choose M_j such that

$$\mathcal{I}_s(M_j) = \inf_g \int_{M_j} |s_g|^2 d\mu_g$$

is realized by an *Einstein* metric g_j with $\lambda < 0$.

Why is Dimension Four Exceptional?

Why is Dimension Four Exceptional?

The Lie group $SO(4)$ is *not simple*:

Why is Dimension Four Exceptional?

The Lie group $SO(4)$ is *not simple*:

$$\mathfrak{so}(4) \cong \mathfrak{so}(3) \oplus \mathfrak{so}(3).$$

Why is Dimension Four Exceptional?

The Lie group $SO(4)$ is *not simple*:

$$\mathfrak{so}(4) \cong \mathfrak{so}(3) \oplus \mathfrak{so}(3).$$

On oriented (M^4, g) ,

Why is Dimension Four Exceptional?

The Lie group $SO(4)$ is *not simple*:

$$\mathfrak{so}(4) \cong \mathfrak{so}(3) \oplus \mathfrak{so}(3).$$

On oriented (M^4, g) , \implies

$$\Lambda^2 = \Lambda^+ \oplus \Lambda^-$$

Why is Dimension Four Exceptional?

The Lie group $SO(4)$ is *not simple*:

$$\mathfrak{so}(4) \cong \mathfrak{so}(3) \oplus \mathfrak{so}(3).$$

On oriented (M^4, g) , \implies

$$\Lambda^2 = \Lambda^+ \oplus \Lambda^-$$

where Λ^\pm are (± 1) -eigenspaces of

$$\star : \Lambda^2 \rightarrow \Lambda^2,$$

$$\star^2 = 1.$$

Why is Dimension Four Exceptional?

The Lie group $SO(4)$ is *not simple*:

$$\mathfrak{so}(4) \cong \mathfrak{so}(3) \oplus \mathfrak{so}(3).$$

On oriented (M^4, g) , \implies

$$\Lambda^2 = \Lambda^+ \oplus \Lambda^-$$

where Λ^\pm are (± 1) -eigenspaces of

$$\star : \Lambda^2 \rightarrow \Lambda^2,$$

$$\star^2 = 1.$$

Λ^+ self-dual 2-forms.

Why is Dimension Four Exceptional?

The Lie group $SO(4)$ is *not simple*:

$$\mathfrak{so}(4) \cong \mathfrak{so}(3) \oplus \mathfrak{so}(3).$$

On oriented (M^4, g) , \implies

$$\Lambda^2 = \Lambda^+ \oplus \Lambda^-$$

where Λ^\pm are (± 1) -eigenspaces of

$$\star : \Lambda^2 \rightarrow \Lambda^2,$$

$$\star^2 = 1.$$

Λ^+ self-dual 2-forms.

Λ^- anti-self-dual 2-forms.

Importance?

Importance?

Curvature is a bundle-value 2-form!

Importance?

Curvature is a bundle-value 2-form!

Vector-bundle-with-connection (E, ∇)

Importance?

Curvature is a bundle-value 2-form!

Vector-bundle-with-connection (E, ∇) over
oriented Riemannian (M^4, g)

Importance?

Curvature is a bundle-value 2-form!

Vector-bundle-with-connection (E, ∇) over oriented Riemannian (M^4, g) has curvature

$$F_{\nabla} = F^+ + F^-$$

Importance?

Curvature is a bundle-value 2-form!

Vector-bundle-with-connection (E, ∇) over oriented Riemannian (M^4, g) has curvature

$$F_{\nabla} = F^+ + F^-$$

where $F^{\pm} \in \Lambda^{\pm} \otimes \text{End}(E)$.

Importance?

Curvature is a bundle-value 2-form!

Vector-bundle-with-connection (E, ∇) over oriented Riemannian (M^4, g) has curvature

$$F_{\nabla} = F^+ + F^-$$

where $F^{\pm} \in \Lambda^{\pm} \otimes \text{End}(E)$.

If $F^- = 0$,

Importance?

Curvature is a bundle-value 2-form!

Vector-bundle-with-connection (E, ∇) over oriented Riemannian (M^4, g) has curvature

$$F_{\nabla} = F^+ + F^-$$

where $F^{\pm} \in \Lambda^{\pm} \otimes \text{End}(E)$.

If $F^- = 0$, so that $F_{\nabla} = F^+$,

Importance?

Curvature is a bundle-value 2-form!

Vector-bundle-with-connection (E, ∇) over oriented Riemannian (M^4, g) has curvature

$$F_{\nabla} = F^+ + F^-$$

where $F^{\pm} \in \Lambda^{\pm} \otimes \text{End}(E)$.

If $F^- = 0$, so that $F_{\nabla} = F^+$,
 ∇ is called self-dual

Importance?

Curvature is a bundle-value 2-form!

Vector-bundle-with-connection (E, ∇) over oriented Riemannian (M^4, g) has curvature

$$F_{\nabla} = F^+ + F^-$$

where $F^{\pm} \in \Lambda^{\pm} \otimes \text{End}(E)$.

If $F^- = 0$, so that $F_{\nabla} = F^+$,
 ∇ is called self-dual (SD).

Importance?

Curvature is a bundle-value 2-form!

Vector-bundle-with-connection (E, ∇) over oriented Riemannian (M^4, g) has curvature

$$F_{\nabla} = F^+ + F^-$$

where $F^{\pm} \in \Lambda^{\pm} \otimes \text{End}(E)$.

If $F^- = 0$, so that $F_{\nabla} = F^+$,
 ∇ is called self-dual (SD).

Proposition. (Λ^+, ∇)

Importance?

Curvature is a bundle-value 2-form!

Vector-bundle-with-connection (E, ∇) over oriented Riemannian (M^4, g) has curvature

$$F_{\nabla} = F^+ + F^-$$

where $F^{\pm} \in \Lambda^{\pm} \otimes \text{End}(E)$.

If $F^- = 0$, so that $F_{\nabla} = F^+$, ∇ is called self-dual (SD).

Proposition. (Λ^+, ∇) is SD

Importance?

Curvature is a bundle-value 2-form!

Vector-bundle-with-connection (E, ∇) over oriented Riemannian (M^4, g) has curvature

$$F_{\nabla} = F^+ + F^-$$

where $F^{\pm} \in \Lambda^{\pm} \otimes \text{End}(E)$.

If $F^- = 0$, so that $F_{\nabla} = F^+$,
 ∇ is called **self-dual (SD)**.

Proposition. (Λ^+, ∇) is **SD** $\iff g$ is *Einstein*.

Importance?

Curvature is a bundle-value 2-form!

Vector-bundle-with-connection (E, ∇) over oriented Riemannian (M^4, g) has curvature

$$F_{\nabla} = F^+ + F^-$$

where $F^{\pm} \in \Lambda^{\pm} \otimes \text{End}(E)$.

If $F^- = 0$, so that $F_{\nabla} = F^+$,
 ∇ is called self-dual (SD).

Proposition. (Λ^+, ∇) is SD $\iff g$ is Einstein.

Donaldson:

Importance?

Curvature is a bundle-value 2-form!

Vector-bundle-with-connection (E, ∇) over oriented Riemannian (M^4, g) has curvature

$$F_{\nabla} = F^+ + F^-$$

where $F^{\pm} \in \Lambda^{\pm} \otimes \text{End}(E)$.

If $F^- = 0$, so that $F_{\nabla} = F^+$,
 ∇ is called self-dual (SD).

Proposition. (Λ^+, ∇) is SD $\iff g$ is Einstein.

Donaldson: moduli spaces of SD connections

Importance?

Curvature is a bundle-value 2-form!

Vector-bundle-with-connection (E, ∇) over oriented Riemannian (M^4, g) has curvature

$$F_{\nabla} = F^+ + F^-$$

where $F^{\pm} \in \Lambda^{\pm} \otimes \text{End}(E)$.

If $F^- = 0$, so that $F_{\nabla} = F^+$,
 ∇ is called self-dual (SD).

Proposition. (Λ^+, ∇) is SD $\iff g$ is Einstein.

Donaldson: moduli spaces of SD connections
 \implies differential topological invariants of M^4 .

Seiberg-Witten Theory:

Seiberg-Witten Theory:

analogous invariants,

Seiberg-Witten Theory:

analogous invariants, related to scalar curvature.

Seiberg-Witten Theory:

analogous invariants, related to scalar curvature.

Given smooth compact oriented 4-manifold M

Seiberg-Witten Theory:

analogous invariants, related to scalar curvature.

Given smooth compact oriented 4-manifold M and
complex line bundle $L \rightarrow M$

Seiberg-Witten Theory:

analogous invariants, related to scalar curvature.

Given smooth compact oriented 4-manifold M and complex line bundle $L \rightarrow M$ such that

$$c_1(L) \equiv w_2 \pmod{2}$$

Seiberg-Witten Theory:

analogous invariants, related to scalar curvature.

Given smooth compact oriented 4-manifold M and complex line bundle $L \rightarrow M$ such that

$$c_1(L) \equiv w_2 \pmod{2}$$

get twisted spin bundles

Seiberg-Witten Theory:

analogous invariants, related to scalar curvature.

Given smooth compact oriented 4-manifold M and complex line bundle $L \rightarrow M$ such that

$$c_1(L) \equiv w_2 \pmod{2}$$

get twisted spin bundles

$$\mathbb{V}_\pm = \mathbb{S}_\pm \otimes L^{1/2}$$

for any metric g on M .

Seiberg-Witten Theory:

analogous invariants, related to scalar curvature.

Given smooth compact oriented 4-manifold M and complex line bundle $L \rightarrow M$ such that

$$c_1(L) \equiv w_2 \pmod{2}$$

get twisted spin bundles

$$V_{\pm} = S_{\pm} \otimes L^{1/2}$$

for any metric g on M . Called spin^c -structure.

Seiberg-Witten Theory:

analogous invariants, related to scalar curvature.

Given smooth compact oriented 4-manifold M and complex line bundle $L \rightarrow M$ such that

$$c_1(L) \equiv w_2 \pmod{2}$$

get twisted spin bundles

$$\mathbb{V}_{\pm} = \mathbb{S}_{\pm} \otimes L^{1/2}$$

for any metric g on M . Called spin^c -structure.

Every unitary connection ϑ on L

Seiberg-Witten Theory:

analogous invariants, related to scalar curvature.

Given smooth compact oriented 4-manifold M and complex line bundle $L \rightarrow M$ such that

$$c_1(L) \equiv w_2 \pmod{2}$$

get twisted spin bundles

$$\mathbb{V}_{\pm} = \mathbb{S}_{\pm} \otimes L^{1/2}$$

for any metric g on M . Called spin^c -structure.

Every unitary connection ϑ on L induces

spin^c Dirac operator

$$D^{\vartheta} : \Gamma(\mathbb{V}_+) \rightarrow \Gamma(\mathbb{V}_-)$$

Seiberg-Witten equations:

$$\begin{aligned} D^{\vartheta} \Phi &= 0 \\ F_{\vartheta}^{+} &= -\frac{1}{2} \Phi \odot \bar{\Phi} \end{aligned}$$

Seiberg-Witten equations:

$$\begin{aligned} D^{\vartheta} \Phi &= 0 \\ F_{\vartheta}^{+} &= -\frac{1}{2} \Phi \odot \bar{\Phi} \end{aligned}$$

Unknowns:

both Φ and ϑ .

Seiberg-Witten equations:

$$\begin{aligned} D^{\mathcal{V}}\Phi &= 0 \\ F_{\mathcal{V}}^+ &= -\frac{1}{2}\Phi \odot \bar{\Phi} \end{aligned}$$

Unknowns:

both Φ and \mathcal{V} .

Here $F_{\mathcal{V}}^+$ = self-dual part of curvature of \mathcal{V} .

Seiberg-Witten equations:

$$\begin{aligned} D^{\mathcal{D}}\Phi &= 0 \\ F_{\mathcal{D}}^+ &= -\frac{1}{2}\Phi \odot \bar{\Phi} \end{aligned}$$

Unknowns:

both Φ and \mathcal{D} .

Here $F_{\mathcal{D}}^+$ = self-dual part of curvature of \mathcal{D} .

Non-linear, but elliptic

Seiberg-Witten equations:

$$\begin{aligned} D^{\vartheta} \Phi &= 0 \\ F_{\vartheta}^{+} &= -\frac{1}{2} \Phi \odot \bar{\Phi} \end{aligned}$$

Unknowns:

both Φ and ϑ .

Here F_{ϑ}^{+} = self-dual part of curvature of ϑ .

Non-linear, but elliptic once ‘gauge-fixing’

$$d^{*}(\vartheta - \vartheta_0) = 0$$

imposed to eliminate automorphisms of $L \rightarrow M$.

Seiberg-Witten equations:

$$\begin{aligned} D^{\mathcal{V}}\Phi &= 0 \\ F_{\mathcal{V}}^+ &= -\frac{1}{2}\Phi \odot \bar{\Phi} \end{aligned}$$

Unknowns:

both Φ and \mathcal{V} .

Here $F_{\mathcal{V}}^+$ = self-dual part of curvature of \mathcal{V} .

Key idea: “counting” solutions defines invariant.

Seiberg-Witten equations:

$$\begin{aligned} D^{\mathcal{V}}\Phi &= 0 \\ F_{\mathcal{V}}^+ &= -\frac{1}{2}\Phi \odot \bar{\Phi} \end{aligned}$$

Unknowns:

both Φ and \mathcal{V} .

Here $F_{\mathcal{V}}^+$ = self-dual part of curvature of \mathcal{V} .

Key idea: “counting” solutions defines invariant.

With mild hypotheses, independent of g .

Seiberg-Witten equations:

$$\begin{aligned} D^{\mathcal{V}}\Phi &= 0 \\ F_{\mathcal{V}}^+ &= -\frac{1}{2}\Phi \odot \bar{\Phi} \end{aligned}$$

Unknowns:

both Φ and \mathcal{V} .

Here $F_{\mathcal{V}}^+$ = self-dual part of curvature of \mathcal{V} .

Key idea: “counting” solutions defines invariant.

With mild hypotheses, independent of g .

Depends only on M & spin^c structure.

Seiberg-Witten equations:

$$\begin{aligned} D^{\vartheta} \Phi &= 0 \\ F_{\vartheta}^{+} &= -\frac{1}{2} \Phi \odot \bar{\Phi} \end{aligned}$$

Unknowns:

both Φ and ϑ .

Here F_{ϑ}^{+} = self-dual part of curvature of ϑ .

Weitzenböck formula:

$$0 = 2\Delta |\Phi|^2 + 4|\nabla^{\vartheta} \Phi|^2 + s|\Phi|^2 + |\Phi|^4$$

Seiberg-Witten equations:

$$\begin{aligned} D^{\mathcal{V}}\Phi &= 0 \\ F_{\mathcal{V}}^+ &= -\frac{1}{2}\Phi \odot \bar{\Phi} \end{aligned}$$

Unknowns:

both Φ and \mathcal{V} .

Here $F_{\mathcal{V}}^+$ = self-dual part of curvature of \mathcal{V} .

Weitzenböck formula:

$$0 = 2\Delta|\Phi|^2 + 4|\nabla^{\mathcal{V}}\Phi|^2 + s|\Phi|^2 + |\Phi|^4$$

gives scalar curvature key role in theory.

Theorem (L '99). *Let M^4 be underlying smooth manifold of a compact complex surface (M, J) with b_1 even.*

Theorem (L '99). *Let M^4 be underlying smooth manifold of a compact complex surface (M, J) with b_1 even. Then $\mathcal{I}_s(M) \neq 0$*

Theorem (L '99). *Let M^4 be underlying smooth manifold of a compact complex surface (M, J) with b_1 even. Then $\mathcal{I}_s(M) \neq 0 \iff$*

Theorem (L '99). *Let M^4 be underlying smooth manifold of a compact complex surface (M, J) with b_1 even. Then $\mathcal{I}_s(M) \neq 0 \iff (M, J)$ is of general type.*

Theorem (L '99). *Let M^4 be underlying smooth manifold of a compact complex surface (M, J) with b_1 even. Then $\mathcal{I}_s(M) \neq 0 \iff (M, J)$ is of general type.*

Recall: $\mathcal{I}_s(M^4) := \inf_g \int_M s_g^2 d\mu_g$

Theorem (L '99). *Let M^4 be underlying smooth manifold of a compact complex surface (M, J) with b_1 even. Then $\mathcal{I}_s(M) \neq 0 \iff (M, J)$ is of general type.*

Recall: $\mathcal{I}_s(M^4) := \inf_g \int_M s_g^2 d\mu_g$

General type means $\dim \Gamma(M, \mathcal{O}(K^{\otimes \ell}))$ grows quadratically as $\ell \rightarrow +\infty$,

Theorem (L '99). *Let M^4 be underlying smooth manifold of a compact complex surface (M, J) with b_1 even. Then $\mathcal{I}_s(M) \neq 0 \iff (M, J)$ is of general type.*

Recall: $\mathcal{I}_s(M^4) := \inf_g \int_M s_g^2 d\mu_g$

General type means $\dim \Gamma(M, \mathcal{O}(K^{\otimes \ell}))$ grows quadratically as $\ell \rightarrow +\infty$, where K denotes the canonical line bundle $\Lambda^{2,0}$.

Theorem (L '99). Let M^4 be underlying smooth manifold of a compact complex surface (M, J) with b_1 even. Then $\mathcal{I}_s(M) \neq 0 \iff (M, J)$ is of general type.

Recall: $\mathcal{I}_s(M^4) := \inf_g \int_M s_g^2 d\mu_g$

General type means $\dim \Gamma(M, \mathcal{O}(K^{\otimes \ell}))$ grows quadratically as $\ell \rightarrow +\infty$, where K denotes the canonical line bundle $\Lambda^{2,0}$.

For complex surfaces, b_1 even \iff Kähler type.

Theorem (L '99). Let M^4 be underlying smooth manifold of a compact complex surface (M, J) with b_1 even. Then $\mathcal{I}_s(M) \neq 0 \iff (M, J)$ is of general type.

Recall: $\mathcal{I}_s(M^4) := \inf_g \int_M s_g^2 d\mu_g$

General type means $\dim \Gamma(M, \mathcal{O}(K^{\otimes \ell}))$ grows quadratically as $\ell \rightarrow +\infty$, where K denotes the canonical line bundle $\Lambda^{2,0}$.


For complex surfaces, b_1 even \iff Kähler type.

Conjecture. For any compact complex surface (M^4, J) with b_1 odd, $\mathcal{I}_s(M) = 0$.

Blowing up:

Blowing up:

If N is a complex surface, may replace $p \in N$
with $\mathbb{C}P_1$




Blowing up:

If N is a complex surface, may replace $p \in N$ with $\mathbb{C}P_1$ to obtain **blow-up**

$$M \approx N \# \overline{\mathbb{C}P_2}$$

in which new $\mathbb{C}P_1$ has self-intersection -1 .



Blowing up:

If N is a complex surface, may replace $p \in N$ with $\mathbb{C}P_1$ to obtain **blow-up**

$$M \approx N \# \overline{\mathbb{C}P_2}$$

in which new $\mathbb{C}P_1$ has self-intersection -1 .

A complex surface X is called **minimal** if it is not the blow-up of another complex surface.

Blowing up:

If N is a complex surface, may replace $p \in N$ with $\mathbb{C}P_1$ to obtain **blow-up**

$$M \approx N \# \overline{\mathbb{C}P_2}$$

in which new $\mathbb{C}P_1$ has self-intersection -1 .

A complex surface X is called **minimal** if it is not the blow-up of another complex surface.

Any complex surface M can be obtained from a minimal surface X by blowing up a finite number of times:

$$M \approx X \# k \overline{\mathbb{C}P_2}$$

One says that X is **minimal model** of M .

Theorem (L '96). *Let (M^4, J) be a compact complex surface of general type,*

Theorem (L '96). *Let (M^4, J) be a compact complex surface of general type, and let X be its minimal model.*

Theorem (L '96). *Let (M^4, J) be a compact complex surface of general type, and let X be its minimal model. Then*

$$\mathcal{I}_s(M) > 0.$$

Theorem (L '96). *Let (M^4, J) be a compact complex surface of general type, and let X be its minimal model. Then*

$$\mathcal{I}_s(M) = 32\pi^2 c_1^2(X) > 0.$$

Theorem (L '96). *Let (M^4, J) be a compact complex surface of general type, and let X be its minimal model. Then*

$$\mathcal{I}_s(M) = 32\pi^2 c_1^2(X) > 0.$$

Seiberg-Witten argument \implies every g satisfies

$$\int_M s_g^2 d\mu_g \geq 32\pi^2 c_1^2(X)$$

Theorem (L '96). *Let (M^4, J) be a compact complex surface of general type, and let X be its minimal model. Then*

$$\mathcal{I}_s(M) = 32\pi^2 c_1^2(X) > 0.$$

Seiberg-Witten argument \implies every g satisfies

$$\int_M s_g^2 d\mu_g \geq 32\pi^2 c_1^2(X)$$

Then must exhibit sequence g_j with

$$\int_M s_{g_j}^2 d\mu_{g_j} \searrow 32\pi^2 c_1^2(X)$$

Key observation:

Key observation: Pluri-canonical model X'
admits orbifold Kähler Einstein metric.

Key observation: Pluri-canonical model X'
admits orbifold Kähler Einstein metric.

Follows from Aubin/Yau because $c_1(X') < 0$.

Key observation: Pluri-canonical model X'
admits orbifold Kähler Einstein metric.

Follows from Aubin/Yau because $c_1(X') < 0$.

$X' = \text{image } M \rightarrow \mathbb{C}\mathbb{P}_N \text{ via linear system } K^{\otimes \ell}.$

Key observation: Pluri-canonical model X'
admits orbifold Kähler Einstein metric.

Follows from Aubin/Yau because $c_1(X') < 0$.

$X' = \text{image } M \rightarrow \mathbb{C}\mathbb{P}_N \text{ via linear system } K^{\otimes \ell}.$

Orbifold, modeled on \mathbb{C}^2/Γ , $\Gamma \subset \mathbf{SU}(2)$.

Key observation: Pluri-canonical model X'
admits orbifold Kähler Einstein metric.

Follows from Aubin/Yau because $c_1(X') < 0$.

$X' = \text{image } M \rightarrow \mathbb{C}\mathbb{P}_N \text{ via linear system } K^{\otimes \ell}$.

Orbifold, modeled on \mathbb{C}^2/Γ , $\Gamma \subset \mathbf{SU}(2)$.

X' obtained from X by collapsing $\mathbb{C}\mathbb{P}_1$'s
contained in regions with $c_1 = 0$.

Key observation: Pluri-canonical model X'
admits orbifold Kähler Einstein metric.

Follows from Aubin/Yau because $c_1(X') < 0$.

$X' = \text{image } M \rightarrow \mathbb{C}\mathbb{P}_N \text{ via linear system } K^{\otimes \ell}.$

Orbifold, modeled on \mathbb{C}^2/Γ , $\Gamma \subset \mathbf{SU}(2)$.

X' obtained from X by collapsing $\mathbb{C}\mathbb{P}_1$'s
contained in regions with $c_1 = 0$.

Strategy: replace neighborhood of each orbifold
point with ALE Ricci-flat manifold.

Given $\Gamma \subset \mathbf{SU}(2)$

Given $\Gamma \subset \mathbf{SU}(2)$ finite subgroup,

Given $\Gamma \subset \mathbf{SU}(2)$ finite subgroup,
the orbifold \mathbb{C}^2/Γ

Given $\Gamma \subset \mathbf{SU}(2)$ finite subgroup,
the orbifold \mathbb{C}^2/Γ can be viewed as
singular complex surface

Given $\Gamma \subset \mathbf{SU}(2)$ finite subgroup,
the orbifold \mathbb{C}^2/Γ can be viewed as
singular complex surface by
choosing generators of Γ -invariant polynomials.

Given $\Gamma \subset \mathbf{SU}(2)$ finite subgroup,
the orbifold \mathbb{C}^2/Γ can be viewed as
singular complex surface $\subset \mathbb{C}^3$ by
choosing 3 generators of Γ -invariant polynomials.

Given $\Gamma \subset \mathbf{SU}(2)$ finite subgroup,
the orbifold \mathbb{C}^2/Γ can be viewed as
singular complex surface $\subset \mathbb{C}^3$ by
choosing 3 generators of Γ -invariant polynomials.

Example. $\begin{bmatrix} e^{2\pi i/m} & \\ & e^{-2\pi i/m} \end{bmatrix} \in \mathbf{SU}(2)$

Given $\Gamma \subset \mathbf{SU}(2)$ finite subgroup,
the orbifold \mathbb{C}^2/Γ can be viewed as
singular complex surface $\subset \mathbb{C}^3$ by
choosing 3 generators of Γ -invariant polynomials.

Example. $\begin{bmatrix} e^{2\pi i/m} & \\ & e^{-2\pi i/m} \end{bmatrix} \in \mathbf{SU}(2)$

generates $\Gamma \cong \mathbb{Z}_m$.

Given $\Gamma \subset \mathbf{SU}(2)$ finite subgroup,
the orbifold \mathbb{C}^2/Γ can be viewed as
singular complex surface $\subset \mathbb{C}^3$ by
choosing 3 generators of Γ -invariant polynomials.

Example. $\begin{bmatrix} e^{2\pi i/m} & \\ & e^{-2\pi i/m} \end{bmatrix} \in \mathbf{SU}(2)$

generates $\Gamma \cong \mathbb{Z}_m$. Setting

$$u = z_1^m, \quad v = z_2^m, \quad y = z_1 z_2,$$

Given $\Gamma \subset \mathbf{SU}(2)$ finite subgroup,
the orbifold \mathbb{C}^2/Γ can be viewed as
singular complex surface $\subset \mathbb{C}^3$ by
choosing 3 generators of Γ -invariant polynomials.

Example. $\begin{bmatrix} e^{2\pi i/m} & \\ & e^{-2\pi i/m} \end{bmatrix} \in \mathbf{SU}(2)$

generates $\Gamma \cong \mathbb{Z}_m$. Setting

$$u = z_1^m, \quad v = z_2^m, \quad y = z_1 z_2,$$

then identifies \mathbb{C}^2/Γ with

$$uv = y^m.$$

Given $\Gamma \subset \mathbf{SU}(2)$ finite subgroup,
the orbifold \mathbb{C}^2/Γ can be viewed as
singular complex surface $\subset \mathbb{C}^3$ by
choosing 3 generators of Γ -invariant polynomials.

Example.
$$\begin{bmatrix} e^{2\pi i/m} & \\ & e^{-2\pi i/m} \end{bmatrix} \in \mathbf{SU}(2)$$

generates $\Gamma \cong \mathbb{Z}_m$. Setting


$$w = \frac{1}{2}(z_1^m - z_2^m), \quad x = \frac{i}{2}(z_1^m + z_2^m), \quad y = z_1 z_2,$$

then identifies \mathbb{C}^2/Γ with

$$w^2 + x^2 + y^m = 0.$$


Felix Klein's Table of Singularities (1884)

Felix Klein's Table of Singularities (1884)


 \mathbb{Z}_m 

$$w^2 + x^2 + y^m = 0$$

Felix Klein's Table of Singularities (1884)


 \mathbb{Z}_m 

$$w^2 + x^2 + y^m = 0$$

 Dih_m^* 

$$w^2 + y(x^2 + y^m) = 0$$


Felix Klein's Table of Singularities (1884)



\mathbb{Z}_m



$$w^2 + x^2 + y^m = 0$$



Dih_m^*



$$w^2 + y(x^2 + y^m) = 0$$




T^*



$$w^2 + x^3 + y^4 = 0$$


Felix Klein's Table of Singularities (1884)



Z_m



$$w^2 + x^2 + y^m = 0$$



Dih_m^*




$$w^2 + y(x^2 + y^m) = 0$$



T^*



$$w^2 + x^3 + y^4 = 0$$




O^*



$$w^2 + x^3 + xy^3 = 0$$


Felix Klein's Table of Singularities (1884)



Z_m



$$w^2 + x^2 + y^m = 0$$



Dih_m^*




$$w^2 + y(x^2 + y^m) = 0$$



T^*




$$w^2 + x^3 + y^4 = 0$$



O^*



$$w^2 + x^3 + xy^3 = 0$$



I^*



$$w^2 + x^3 + y^5 = 0$$

Prototypical Klein singularity:

$$w^2 + x^2 + y^2 = 0$$

Two ways to get rid of a singularity:

$$w^2 + x^2 + y^2 = 0$$

Two ways to get rid of a singularity:


- Smooth it, by deformation:


$$w^2 + x^2 + y^2 = 0$$


Two ways to get rid of a singularity:


- Smooth it, by deformation:


$$w^2 + x^2 + y^2 = \epsilon$$














Two ways to get rid of a singularity:

- Smooth it, by deformation:

$$w^2 + x^2 + y^2 = \epsilon$$

Two ways to get rid of a singularity:

- Smooth it, by deformation:

$$w^2 + x^2 + y^2 = \epsilon$$

- Resolve it, by blowing up,

$$w^2 + x^2 + y^2 = 0$$

Two ways to get rid of a singularity:

- Smooth it, by deformation:

$$w^2 + x^2 + y^2 = \epsilon$$

- Resolve it, by blowing up,

$$w^2 + x^2 + y^2 = 0$$

$$\mathcal{O}(-1)$$



$$\mathbb{C}P_2$$

Two ways to get rid of a singularity:

- Smooth it, by deformation:

$$w^2 + x^2 + y^2 = \epsilon$$

- Resolve it, by blowing up,

$$w^2 + x^2 + y^2 = 0$$

$$\begin{array}{ccc} & & \mathcal{O}(-1) \\ & & \downarrow \\ \mathbb{CP}_1 & \hookrightarrow & \mathbb{CP}_2 \end{array}$$

Two ways to get rid of a singularity:


- Smooth it, by deformation:


$$w^2 + x^2 + y^2 = \epsilon$$


- Resolve it, by blowing up, iteratively:


$$w^2 + x^2 + y^2 = 0$$


$$\begin{array}{ccc} \mathcal{O}(-2) & \rightarrow & \mathcal{O}(-1) \\ \downarrow & & \downarrow \\ \mathbb{C}P_1 & \hookrightarrow & \mathbb{C}P_2 \end{array}$$














Two ways to get rid of a singularity:

- Smooth it, by deformation:

$$w^2 + x^2 + y^2 = \epsilon$$

- Resolve it, by blowing up, iteratively:

$$w^2 + x^2 + y^2 = 0$$

$$\begin{array}{ccc} \mathcal{O}(-2) & \rightarrow & \mathcal{O}(-1) \\ \downarrow & & \downarrow \\ \mathbb{C}P_1 & \hookrightarrow & \mathbb{C}P_2 \end{array}$$

Two ways to get rid of a singularity:

- Smooth it, by deformation:

$$w^2 + x^2 + y^2 = \epsilon$$

- Resolve it, by blowing up, iteratively:

$$w^2 + x^2 + y^2 = 0$$

$$\begin{array}{ccc} \mathcal{O}(-2) & \rightarrow & \mathcal{O}(-1) \\ \downarrow & & \downarrow \\ \mathbb{C}P_1 & \hookrightarrow & \mathbb{C}P_2 \end{array}$$

Usually these are topologically different.

Two ways to get rid of a singularity:

- Smooth it, by deformation:

$$w^2 + x^2 + y^2 = \epsilon$$

- Resolve it, by blowing up, iteratively:

$$w^2 + x^2 + y^2 = 0$$

$$\begin{array}{ccc} \mathcal{O}(-2) & \rightarrow & \mathcal{O}(-1) \\ \downarrow & & \downarrow \\ \mathbb{C}P_1 & \hookrightarrow & \mathbb{C}P_2 \end{array}$$

Usually these are topologically different.

But for Klein singularities, they are diffeomorphic!

Two ways to get rid of a singularity:

- Smooth it, by deformation:

$$w^2 + x^2 + y^2 = \epsilon$$

- Resolve it, by blowing up, iteratively:

$$w^2 + x^2 + y^2 = 0$$

$$\begin{array}{ccc} \mathcal{O}(-2) & \rightarrow & \mathcal{O}(-1) \\ \downarrow & & \downarrow \\ \mathbb{CP}_1 & \hookrightarrow & \mathbb{CP}_2 \end{array}$$

Usually these are topologically different.

But for Klein singularities, they are diffeomorphic!

Gorenstein singularities.

Two ways to get rid of a singularity:

- Smooth it, by deformation:

$$w^2 + x^2 + y^2 = \epsilon$$

- Resolve it, by blowing up, iteratively:

$$w^2 + x^2 + y^2 = 0$$

$$\begin{array}{ccc} \mathcal{O}(-2) & \rightarrow & \mathcal{O}(-1) \\ \downarrow & & \downarrow \\ \mathbb{C}P_1 & \hookrightarrow & \mathbb{C}P_2 \end{array}$$

Usually these are topologically different.

But for Klein singularities, they are diffeomorphic!

Gorenstein singularities. Crepant Resolutions.

Resolutions of Klein Singularities:

Resolutions of Klein Singularities:

\forall Klein singularity $V \subset \mathbb{C}^3$,

Resolutions of Klein Singularities:

\forall Klein singularity $V \subset \mathbb{C}^3$, $\exists!$ resolution

$$\hat{V} \rightarrow V$$

Resolutions of Klein Singularities:

\forall Klein singularity $V \subset \mathbb{C}^3$, $\exists!$ resolution

$$\hat{V} \rightarrow V$$

with $c_1(T^{1,0}\hat{V}) = 0$.


Resolutions of Klein Singularities:

\forall Klein singularity $V \subset \mathbb{C}^3$, $\exists!$ resolution

$$\hat{V} \rightarrow V$$

with $c_1(T^{1,0}\hat{V}) = 0$.

Replaces origin with a union of $\mathbb{C}P_1$'s,




Resolutions of Klein Singularities:

\forall Klein singularity $V \subset \mathbb{C}^3$, $\exists!$ resolution

$$\hat{V} \rightarrow V$$

with $c_1(T^{1,0}\hat{V}) = 0$.

Replaces origin with a union of $\mathbb{C}P_1$'s,
each with self-intersection -2 ,




Resolutions of Klein Singularities:

\forall Klein singularity $V \subset \mathbb{C}^3$, $\exists!$ resolution

$$\hat{V} \rightarrow V$$

with $c_1(T^{1,0}\hat{V}) = 0$.

Replaces origin with a union of $\mathbb{C}P_1$'s,
each with self-intersection -2 ,
meeting transversely,




Resolutions of Klein Singularities:

\forall Klein singularity $V \subset \mathbb{C}^3$, $\exists!$ resolution

$$\hat{V} \rightarrow V$$

with $c_1(T^{1,0}\hat{V}) = 0$.

Replaces origin with a union of $\mathbb{C}P_1$'s,
each with self-intersection -2 ,
meeting transversely, & forming connected set:




Resolutions of Klein Singularities:

\forall Klein singularity $V \subset \mathbb{C}^3$, $\exists!$ resolution


$$\hat{V} \rightarrow V$$

with $c_1(T^{1,0}\hat{V}) = 0$.

Replaces origin with a union of $\mathbb{C}P_1$'s,
each with self-intersection -2 ,
meeting transversely, & forming connected set:



Intersection pattern dual to Dynkin diagram!




Resolutions of Klein Singularities:

\forall Klein singularity $V \subset \mathbb{C}^3$, $\exists!$ resolution


$$\hat{V} \rightarrow V$$

with $c_1(T^{1,0}\hat{V}) = 0$.

Replaces origin with a union of $\mathbb{C}\mathbb{P}_1$'s,
each with self-intersection -2 ,
meeting transversely, & forming connected set:



Intersection pattern dual to Dynkin diagram!




Resolutions of Klein Singularities:

\forall Klein singularity $V \subset \mathbb{C}^3$, $\exists!$ resolution


$$\hat{V} \rightarrow V$$

with $c_1(T^{1,0}\hat{V}) = 0$.

Replaces origin with a union of $\mathbb{C}P_1$'s,
each with self-intersection -2 ,
meeting transversely, & forming connected set:



Intersection pattern dual to Dynkin diagram!




Resolutions of Klein Singularities:

\forall Klein singularity $V \subset \mathbb{C}^3$, $\exists!$ resolution


$$\hat{V} \rightarrow V$$

with $c_1(T^{1,0}\hat{V}) = 0$.

Replaces origin with a union of \mathbb{CP}_1 's,
each with self-intersection -2 ,
meeting transversely, & forming connected set:



Intersection pattern dual to Dynkin diagram!




Resolutions of Klein Singularities:

\forall Klein singularity $V \subset \mathbb{C}^3$, $\exists!$ resolution


$$\hat{V} \rightarrow V$$


with $c_1(T^{1,0}\hat{V}) = 0$.

Replaces origin with a union of $\mathbb{C}P_1$'s,
each with self-intersection -2 ,
meeting transversely, & forming connected set:




Intersection pattern dual to Dynkin diagram!






$$\mathbb{Z}_{k+1} \longleftrightarrow A_k$$







$$\mathbb{Z}_{k+1} \longleftrightarrow A_k$$




$$\text{Dih}_{k-2}^* \longleftrightarrow D_k$$







$$\mathbb{Z}_{k+1} \longleftrightarrow A_k$$




$$\text{Dih}_{k-2}^* \longleftrightarrow D_k$$




$$T^* \longleftrightarrow E_6$$








$$\mathbb{Z}_{k+1} \longleftrightarrow A_k$$




$$\text{Dih}_{k-2}^* \longleftrightarrow D_k$$




$$T^* \longleftrightarrow E_6$$




$$O^* \longleftrightarrow E_7$$








$$\mathbb{Z}_{k+1} \longleftrightarrow A_k$$





$$\text{Dih}_{k-2}^* \longleftrightarrow D_k$$



$$T^* \longleftrightarrow E_6$$



$$O^* \longleftrightarrow E_7$$



$$I^* \longleftrightarrow E_8$$



McKay Correspondence

McKay Correspondence

Given $\rho : \Gamma \hookrightarrow \mathbf{SU}(2)$ finite subgroup,

McKay Correspondence

Given $\rho : \Gamma \hookrightarrow \mathbf{SU}(2)$ finite subgroup,
construct Dynkin diagram, as follows:

McKay Correspondence

Given $\rho : \Gamma \hookrightarrow \mathbf{SU}(2)$ finite subgroup,
construct Dynkin diagram, as follows:

One node for each non-trivial irred. representation

$$\rho_j : \Gamma \rightarrow \text{End}(\mathbb{V}_j)$$

McKay Correspondence

Given $\rho : \Gamma \hookrightarrow \mathbf{SU}(2)$ finite subgroup,
construct Dynkin diagram, as follows:

One node for each non-trivial irred. representation

$$\rho_j : \Gamma \rightarrow \text{End}(\mathbb{V}_j)$$

Next decompose

$$\rho \otimes \rho_j = \bigoplus_{\ell} (\rho_{\ell})^{\oplus n_{j\ell}}$$

McKay Correspondence

Given $\rho : \Gamma \hookrightarrow \mathbf{SU}(2)$ finite subgroup,
construct Dynkin diagram, as follows:

One node for each non-trivial irred. representation

$$\rho_j : \Gamma \rightarrow \text{End}(\mathbb{V}_j)$$

Next decompose

$$\rho \otimes \rho_j = \bigoplus_{\ell} (\rho_{\ell})^{\oplus n_{j\ell}}$$

as sum of irreducibles. Then $n_{j\ell} = n_{\ell j} = 0$ or 1 .

McKay Correspondence

Given $\rho : \Gamma \hookrightarrow \mathbf{SU}(2)$ finite subgroup,
construct Dynkin diagram, as follows:

One node for each non-trivial irred. representation

$$\rho_j : \Gamma \rightarrow \text{End}(\mathbb{V}_j)$$

Next decompose

$$\rho \otimes \rho_j = \bigoplus_{\ell} (\rho_{\ell})^{\oplus n_{j\ell}}$$

as sum of irreducibles. Then $n_{j\ell} = n_{\ell j} = 0$ or 1 .

Now draw edge joining nodes j & ℓ if $n_{j\ell} \neq 0$.

McKay Correspondence

Given $\rho : \Gamma \hookrightarrow \mathbf{SU}(2)$ finite subgroup,
construct Dynkin diagram, as follows:

One node for each **non-trivial** irred. representation

$$\rho_j : \Gamma \rightarrow \text{End}(\mathbb{V}_j)$$


Next decompose

$$\rho \otimes \rho_j = \bigoplus_{\ell} (\rho_{\ell})^{\oplus n_{j\ell}}$$


as sum of irreducibles. Then $n_{j\ell} = n_{\ell j} = 0$ or 1 .

Now draw **edge** joining nodes j & ℓ if $n_{j\ell} \neq 0$.


Reproduces Dynkin diagram of crepant resolution!





$$\mathbb{Z}_{k+1} \longleftrightarrow A_k$$





$$\text{Dih}_{k-2}^* \longleftrightarrow D_k$$




$$T^* \longleftrightarrow E_6$$



$$O^* \longleftrightarrow E_7$$



$$I^* \longleftrightarrow E_8$$



Theorem (Kronheimer '89). *Let $\Gamma \subset \mathbf{SU}(2)$ finite, and let \mathbf{Y} be the smooth 4-manifold gotten by resolving the singularity of \mathbb{R}^4/Γ .*

Theorem (Kronheimer '89). *Let $\Gamma \subset \mathbf{SU}(2)$ finite, and let \mathbf{Y} be the smooth 4-manifold gotten by resolving the singularity of \mathbb{R}^4/Γ . Then \mathbf{Y} admits a family of *ALE Ricci-flat metrics*.*


Theorem (Kronheimer '89). *Let $\Gamma \subset \mathbf{SU}(2)$ finite, and let \mathbf{Y} be the smooth 4-manifold gotten by resolving the singularity of \mathbb{R}^4/Γ . Then \mathbf{Y} admits a family of *ALE Ricci-flat metrics*. Every such metric is *hyper-Kähler*,*

(M^n, g) :

holonomy


(M^n, g) :

holonomy




(M^n, g) :

holonomy




(M^n, g) :

holonomy




(M^n, g) :

holonomy




(M^n, g) :

holonomy




(M^n, g) :

holonomy




(M^n, g) :

holonomy




(M^n, g) :

holonomy




(M^n, g) :

holonomy




(M^n, g) :

holonomy




(M^n, g) :

holonomy




(M^n, g) :

holonomy




(M^n, g) :

holonomy




(M^n, g) :

holonomy




(M^n, g) :

holonomy



$(M^n, g):$


holonomy $\subset O(n)$



Kähler metrics:


(M^{2m}, g) :

holonomy




Kähler metrics:

(M^{2m}, g) Kähler \iff holonomy $\subset \mathbf{U}(m)$




Kähler metrics:

(M^4, g) Kähler \iff holonomy $\subset \mathbf{U}(2)$




Hyper-Kähler metrics:

$(M^{4\ell}, g)$ Hyper-Kähler \iff holonomy $\subset \mathbf{Sp}(\ell)$




Hyper-Kähler metrics:

(M^4, g) Hyper-Kähler \iff holonomy $\subset \mathbf{Sp}(1)$



Hyper-Kähler metrics:

(M^4, g) Hyper-Kähler \iff holonomy $\subset \mathbf{SU}(2)$



Theorem (Kronheimer '89). *Let $\Gamma \subset \mathbf{SU}(2)$ finite, and let \mathbf{Y} be the smooth 4-manifold gotten by resolving the singularity of \mathbb{R}^4/Γ . Then \mathbf{Y} admits a family of *ALE Ricci-flat metrics*. Every such metric is *hyper-Kähler*,*

Theorem (Kronheimer '89). *Let $\Gamma \subset \mathbf{SU}(2)$ finite, and let \mathbf{Y} be the smooth 4-manifold gotten by resolving the singularity of \mathbb{R}^4/Γ . Then \mathbf{Y} admits a family of *ALE Ricci-flat metrics*. Every such metric is *hyper-Kähler*, and the moduli space of these metrics is connected:*

Theorem (Kronheimer '89). *Let $\Gamma \subset \mathbf{SU}(2)$ finite, and let \mathbf{Y} be the smooth 4-manifold gotten by resolving the singularity of \mathbb{R}^4/Γ . Then \mathbf{Y} admits a family of *ALE Ricci-flat metrics*. Every such metric is *hyper-Kähler*, and the moduli space of these metrics is connected: roughly, $[\text{Weyl chamber}]^3 \subset \mathfrak{h} \otimes \mathbb{R}^3$, where $\mathfrak{h} \cong H^2(\mathbf{Y}, \mathbb{R})$ is Cartan of Lie algebra \mathfrak{g} corresponding to Dynkin diagram associated with Γ .*

Theorem (Kronheimer '89). *Let $\Gamma \subset \mathbf{SU}(2)$ finite, and let \mathbf{Y} be the smooth 4-manifold gotten by resolving the singularity of \mathbb{R}^4/Γ . Then \mathbf{Y} admits a family of *ALE Ricci-flat metrics*. Every such metric is *hyper-Kähler*, and the moduli space of these metrics is connected: roughly, $[\text{Weyl chamber}]^3 \subset \mathfrak{h} \otimes \mathbb{R}^3$, where $\mathfrak{h} \cong H^2(\mathbf{Y}, \mathbb{R})$ is Cartan of Lie algebra \mathfrak{g} corresponding to Dynkin diagram associated with Γ .*

A_1 : Eguchi-Hanson

Theorem (Kronheimer '89). *Let $\Gamma \subset \mathbf{SU}(2)$ finite, and let \mathbf{Y} be the smooth 4-manifold gotten by resolving the singularity of \mathbb{R}^4/Γ . Then \mathbf{Y} admits a family of *ALE Ricci-flat metrics*. Every such metric is *hyper-Kähler*, and the moduli space of these metrics is connected: roughly, $[\text{Weyl chamber}]^3 \subset \mathfrak{h} \otimes \mathbb{R}^3$, where $\mathfrak{h} \cong H^2(\mathbf{Y}, \mathbb{R})$ is Cartan of Lie algebra \mathfrak{g} corresponding to Dynkin diagram associated with Γ .*

A_1 : Eguchi-Hanson

A_k : Gibbons-Hawking/Hitchin

Theorem (Kronheimer '89). *Let $\Gamma \subset \mathbf{SU}(2)$ finite, and let \mathbf{Y} be the smooth 4-manifold gotten by resolving the singularity of \mathbb{R}^4/Γ . Then \mathbf{Y} admits a family of *ALE Ricci-flat metrics*. Every such metric is *hyper-Kähler*, and the moduli space of these metrics is connected: roughly, $[\text{Weyl chamber}]^3 \subset \mathfrak{h} \otimes \mathbb{R}^3$, where $\mathfrak{h} \cong H^2(\mathbf{Y}, \mathbb{R})$ is Cartan of Lie algebra \mathfrak{g} corresponding to Dynkin diagram associated with Γ .*

A_1 : Eguchi-Hanson

A_k : Gibbons-Hawking/Hitchin

D_k : conjectured by Hitchin

Hyper-Kähler quotient (HKLR)

Hyper-Kähler quotient (HKLR)

If G compact acts on simply-connected
hyper-Kähler \mathbf{V}

Hyper-Kähler quotient (HKLR)

If G compact acts on simply-connected
hyper-Kähler \mathbf{V} by tri-holomorphic isometries,

Hyper-Kähler quotient (HKLR)

If G compact acts on simply-connected
hyper-Kähler \mathbf{V} by tri-holomorphic isometries,
moment map is G -equivariant map

Hyper-Kähler quotient (HKLR)

If G compact acts on simply-connected
hyper-Kähler \mathbf{V} by tri-holomorphic isometries,
moment map is G -equivariant map

$$\mu : \mathbf{V} \rightarrow \mathfrak{g}^* \otimes \mathbb{R}^3$$

Hyper-Kähler quotient (HKLR)

If G compact acts on simply-connected
hyper-Kähler \mathbf{V} by tri-holomorphic isometries,
moment map is G -equivariant map

$$\mu : \mathbf{V} \rightarrow \mathfrak{g}^* \otimes \mathbb{R}^3$$

encoding Hamiltonians for generators.

Hyper-Kähler quotient (HKLR)

If G compact acts on simply-connected hyper-Kähler \mathbf{V} by tri-holomorphic isometries, moment map is G -equivariant map

$$\mu : \mathbf{V} \rightarrow \mathfrak{g}^* \otimes \mathbb{R}^3$$

encoding Hamiltonians for generators.

If G acts freely on $\mu^{-1}(\zeta)$,

Hyper-Kähler quotient (HKLR)

If G compact acts on simply-connected hyper-Kähler \mathbf{V} by tri-holomorphic isometries, moment map is G -equivariant map

$$\mu : \mathbf{V} \rightarrow \mathfrak{g}^* \otimes \mathbb{R}^3$$

encoding Hamiltonians for generators.

If G acts freely on $\mu^{-1}(\zeta)$, then

$$\mathbf{V} // G := [\mu^{-1}(\zeta)] / G$$

Hyper-Kähler quotient (HKLR)

If G compact acts on simply-connected hyper-Kähler \mathbf{V} by tri-holomorphic isometries, moment map is G -equivariant map

$$\mu : \mathbf{V} \rightarrow \mathfrak{g}^* \otimes \mathbb{R}^3$$

encoding Hamiltonians for generators.

If G acts freely on $\mu^{-1}(\zeta)$, then

$$\mathbf{V} // G := [\mu^{-1}(\zeta)] / G$$

inherits natural hyper-Kähler structure,

Hyper-Kähler quotient (HKLR)

If G compact acts on simply-connected hyper-Kähler \mathbf{V} by tri-holomorphic isometries, moment map is G -equivariant map

$$\mu : \mathbf{V} \rightarrow \mathfrak{g}^* \otimes \mathbb{R}^3$$

encoding Hamiltonians for generators.

If G acts freely on $\mu^{-1}(\zeta)$, then

$$\mathbf{V} // G := [\mu^{-1}(\zeta)] / G$$

inherits natural hyper-Kähler structure, and

$$\dim(\mathbf{V} // G) = \dim \mathbf{V} - 4 \dim G$$

Kronheimer Construction

Kronheimer Construction

$\Gamma \subset \mathbf{SU}(2)$ acts on \mathbb{H} and $\mathbb{R}\Gamma = \mathbb{R}^{|\Gamma|}$.

Kronheimer Construction

$\Gamma \subset \mathbf{SU}(2)$ acts on \mathbb{H} and $\mathbb{R}\Gamma = \mathbb{R}^{|\Gamma|}$. Set

$$\mathbf{V} = [\mathbb{H} \otimes_{\mathbb{R}} \text{End}(\mathbb{R}\Gamma)]^{\Gamma} = [\mathbb{C}^2 \otimes_{\mathbb{C}} \text{End}(\mathbb{C}\Gamma)]^{\Gamma}$$

Kronheimer Construction

$\Gamma \subset \mathbf{SU}(2)$ acts on \mathbb{H} and $\mathbb{R}\Gamma = \mathbb{R}^{|\Gamma|}$. Set

$$\mathbf{V} = [\mathbb{H} \otimes_{\mathbb{R}} \text{End}(\mathbb{R}\Gamma)]^{\Gamma} = [\mathbb{C}^2 \otimes_{\mathbb{C}} \text{End}(\mathbb{C}\Gamma)]^{\Gamma}$$

$$G = [\mathbf{SU}(\mathbb{C}\Gamma)]^{\Gamma} \cong \mathbf{U}(n_1) \times \cdots \times \mathbf{U}(n_k)$$

Kronheimer Construction

$\Gamma \subset \mathbf{SU}(2)$ acts on \mathbb{H} and $\mathbb{R}\Gamma = \mathbb{R}^{|\Gamma|}$. Set

$$\mathbf{V} = [\mathbb{H} \otimes_{\mathbb{R}} \text{End}(\mathbb{R}\Gamma)]^{\Gamma} = [\mathbb{C}^2 \otimes_{\mathbb{C}} \text{End}(\mathbb{C}\Gamma)]^{\Gamma}$$

$$G = [\mathbf{SU}(\mathbb{C}\Gamma)]^{\Gamma} \cong \mathbf{U}(n_1) \times \cdots \times \mathbf{U}(n_k)$$

where n_1, \dots, n_k dim's non-triv. irred. reps. Γ .

Kronheimer Construction

$\Gamma \subset \mathbf{SU}(2)$ acts on \mathbb{H} and $\mathbb{R}\Gamma = \mathbb{R}^{|\Gamma|}$. Set

$$\mathbf{V} = [\mathbb{H} \otimes_{\mathbb{R}} \text{End}(\mathbb{R}\Gamma)]^{\Gamma} = [\mathbb{C}^2 \otimes_{\mathbb{C}} \text{End}(\mathbb{C}\Gamma)]^{\Gamma}$$

$$G = [\mathbf{SU}(\mathbb{C}\Gamma)]^{\Gamma} \cong \mathbf{U}(n_1) \times \cdots \times \mathbf{U}(n_k)$$

where n_1, \dots, n_k dim's non-triv. irred. reps. Γ .

Then $\mathbf{Y} = \mathbf{V} // G$ gives ALE hyper-Kähler structure to resolution of \mathbb{C}^2/Γ .

Kronheimer Construction

$\Gamma \subset \mathbf{SU}(2)$ acts on \mathbb{H} and $\mathbb{R}\Gamma = \mathbb{R}^{|\Gamma|}$. Set

$$\mathbf{V} = [\mathbb{H} \otimes_{\mathbb{R}} \text{End}(\mathbb{R}\Gamma)]^{\Gamma} = [\mathbb{C}^2 \otimes_{\mathbb{C}} \text{End}(\mathbb{C}\Gamma)]^{\Gamma}$$

$$G = [\mathbf{SU}(\mathbb{C}\Gamma)]^{\Gamma} \cong \mathbf{U}(n_1) \times \cdots \times \mathbf{U}(n_k)$$

where n_1, \dots, n_k dim's non-triv. irred. reps. Γ .

Then $\mathbf{Y} = \mathbf{V} // G$ gives ALE hyper-Kähler structure to resolution of \mathbb{C}^2/Γ .

Construction depends on ζ : $3k$ parameters.

Kronheimer's ALE spaces are Ricci-flat and Kähler.

Kronheimer's **ALE** spaces are Ricci-flat and Kähler.

Play ubiquitous role for Einstein 4-manifolds.

Kronheimer's **ALE** spaces are Ricci-flat and **Kähler**.

Play ubiquitous role for Einstein 4-manifolds.

But turns out that **scalar-flat Kähler ALE** spaces also play an important role.

Kronheimer's **ALE** spaces are Ricci-flat and **Kähler**.

Play ubiquitous role for Einstein 4-manifolds.

But turns out that **scalar-flat Kähler ALE** spaces also play an important role.

For example, proof of main result also needs one such example (not discussed here).

Kronheimer's **ALE** spaces are Ricci-flat and **Kähler**.

Play ubiquitous role for Einstein 4-manifolds.

But turns out that **scalar-flat Kähler ALE** spaces also play an important role.

For example, proof of main result also needs one such example (not discussed here).

Also play key role in recent existence proofs for Einstein metrics (**Chen-L-Weber '08, L '12**)

Kronheimer's **ALE** spaces are Ricci-flat and **Kähler**.

Play ubiquitous role for Einstein 4-manifolds.

But turns out that **scalar-flat Kähler ALE** spaces also play an important role.

For example, proof of main result also needs one such example (not discussed here).

Also play key role in recent existence proofs for Einstein metrics (**Chen-L-Weber '08, L '12**)

Problem: Classify them!

Kronheimer's **ALE** spaces are Ricci-flat and **Kähler**.

Play ubiquitous role for Einstein 4-manifolds.

But turns out that **scalar-flat Kähler ALE** spaces also play an important role.

For example, proof of main result also needs one such example (not discussed here).

Also play key role in recent existence proofs for Einstein metrics (**Chen-L-Weber '08, L '12**)

Problem: Classify them!

We know many examples (**L '88, '91, C-S '05**),

Kronheimer's **ALE** spaces are Ricci-flat and **Kähler**.

Play ubiquitous role for Einstein 4-manifolds.

But turns out that **scalar-flat Kähler ALE** spaces also play an important role.

For example, proof of main result also needs one such example (not discussed here).

Also play key role in recent existence proofs for Einstein metrics (**Chen-L-Weber '08, L '12**)

Problem: Classify them!

We know many examples (**L '88, '91, C-S '05**), but have classification only in toric case.

Kronheimer's **ALE** spaces are Ricci-flat and **Kähler**.

Play ubiquitous role for Einstein 4-manifolds.

But turns out that **scalar-flat Kähler ALE** spaces also play an important role.

For example, proof of main result also needs one such example (not discussed here).

Also play key role in recent existence proofs for Einstein metrics (**Chen-L-Weber '08, L '12**)

Problem: Classify them!

We know many examples (**L '88, '91, C-S '05**), but have classification only in toric case.

Generalized McKay correspondence **suggests** approach,

Kronheimer's **ALE** spaces are Ricci-flat and **Kähler**.

Play ubiquitous role for Einstein 4-manifolds.

But turns out that **scalar-flat Kähler ALE** spaces also play an important role.

For example, proof of main result also needs one such example (not discussed here).

Also play key role in recent existence proofs for Einstein metrics (**Chen-L-Weber '08, L '12**)

Problem: Classify them!

We know many examples (**L '88, '91, C-S '05**), but have classification only in toric case.

Generalized McKay correspondence suggests approach, currently under investigation.