

# Non Locally-Connected Julia Sets constructed by iterated tuning

John Milnor

*Stony Brook University*

Revised May 26, 2006

## Notations:

Every quadratic map

$$f_c(z) = z^2 + c$$

has two fixed points,  $\alpha$  and  $\beta$ , where

$$\alpha + \beta = 1, \quad \alpha\beta = c, \quad \Re(\alpha) \leq \Re(\beta).$$

The multiplier

$$\mu = f'_c(\alpha) = 2\alpha$$

will be used as an alternative parameter for the quadratic family.


Here  $c$  and  $\mu$  determine each other:

$$c = c(\mu) = \alpha(1 - \alpha) \quad \text{with} \quad \alpha = \mu/2,$$

$$\mu = \mu(c) = 1 - \sqrt{1 - 4c}, \quad \text{with} \quad \Re(\mu) \leq 1.$$


The map  $f_{c(\mu)}$  corresponding to  $\mu$  will be denoted by

$$\widehat{f}_\mu(z) = z^2 + c(\mu).$$




The **connectedness locus**  $\widehat{M}$ , consisting of all  $\mu$  in the half-plane  $\Re(\mu) \leq 1$  with  $K(\widehat{f}_\mu)$  connected, will be called the **rounded Mandelbrot set**.

*Its **period one hyperbolic component**, the set of all  $\mu \in \widehat{M}$  for which  $\widehat{f}_\mu$  has an attracting fixed point, is the open unit disk  $\mathbb{D}$ .*



Region in  $\hat{M}$



and

Disk Approximation

There is a **satellite** hyperbolic component  $H(n/p)$  of period  $p$  attached to  $\mathbb{D}$  at each  $p$ -th root of unity  $e^{2\pi i n/p}$ .

*Similarly, there are satellites  $H(n/p) \triangleright H(n'/p')$  of period  $pp'$  attached to  $H(n/p)$  at corresponding boundary points; and so on.*

Empirically, each iterated satellite  $H(n_1/p_1) \triangleright \dots \triangleright H(n_k/p_k)$  can be approximated by a round disk of radius  $1/(p_1 \cdots p_k)^2$ .

**Question: How can this be made precise?**

## Tuning (in parameter space).


The Douady-Hubbard **tuning** construction assigns to each hyperbolic component  $H \subset \widehat{M}$  a homeomorphism

$$H \triangleright : \widehat{M} \xrightarrow{\cong} (H \triangleright \widehat{M}) \subset \widehat{M}$$


from  $\widehat{M}$  onto a “small copy” of  $\widehat{M}$ .

- Each  $H \triangleright$  maps hyperbolic components to hyperbolic components, with  $\text{per}(H \triangleright H') = \text{per}(H) \text{per}(H')$ .
- The set of all  $H$  forms a free non-commutative monoid, with  $\mathbb{D}$  as identity element.
- Each  $H \triangleright : \overline{\mathbb{D}} \rightarrow \overline{H}$  is holomorphic, and yields the canonical **Douady-Hubbard parametrization** of  $H$ : For each  $\mu \in \mathbb{D}$ , the attracting periodic orbit for the map  $\widehat{f}_{H \triangleright \mu}$  has multiplier  $\mu$ .
- The image  $H \triangleright 1 \in \partial H$  is called the **root point** of  $H$ .

## Tuning in the dynamic plane (intuitive picture).




$\triangleright$  — =



To obtain the filled Julia set for  $H_1 \triangleright \mu_2$ , choose  $\mu_1 \in H_1$ , then replace every Fatou component of  $K(\widehat{f}_{\mu_1})$  by a copy of  $K(\widehat{f}_{\mu_2})$ . (In the figure,  $\widehat{f}_{\mu_2}$  is the Chebyshev map  $z \mapsto z^2 - 2$ , and  $K(\widehat{f}_{\mu_2})$  is the line segment  $[-2, 2]$ .)

# Constructing a non locally-connected $K(\widehat{f}_\mu)$



(Small disk sizes exaggerated.)

Choose any sequence of rational angles  $t_1, t_2, \dots \neq 0$ ,  
and let  $r_k$  be the root point of

$$H(t_1) \triangleright \dots \triangleright H(t_k).$$

# The Theorem.

Let  $\omega \in \widehat{M}$  be any limit point for the sequence of root points


$$r_k \in \partial\left(H(t_1) \triangleright \cdots \triangleright H(t_k)\right) \quad \text{as} \quad k \rightarrow \infty.$$

## Theorem (Douady, Hubbard, Sørensen).

*If the sequence  $\{|t_j|\}$  converges to zero sufficiently rapidly, then the filled Julia set  $K(\widehat{f}_\omega)$  is not locally-connected.*

The proof will be based on **external rays** and **separating periodic orbits**.






Two nested wakes  $W_{[1/7, 2/7]} \supset W_{[1/5, 4/15]}$ .

Every root point  $H \triangleright 1 \in \widehat{M}$  is the landing point of exactly two external rays, with angles  $0 \leq a < b \leq 1$ . These rays cut the parameter plane into two halves.


**Definition.** The half containing  $H = H_{[a,b]}$  is called the **wake**  $W_{[a,b]}$ , and  $[a, b]$  is called its **characteristic interval**.

## In the Dynamic Plane:



For every hyperbolic component  $H = H_{[a,b]}$  of period  $p > 1$  and every  $\mu \in H$ , the external rays of angle  $a$  and  $b$  for  $K(\hat{f}_\mu)$  land at a common repelling periodic point. I will write


$$z^1 = z^1([a, b], \mu) = \ell_a(\mu) = \ell_b(\mu) \in \partial K(\hat{f}_\mu). \quad 10$$



Number the periodic Fatou components of  $\widehat{f}_\mu$  as  $0 \in U_0 \mapsto U_1 \mapsto \dots$ . Then the orbit of  $z^1$  consists of points


$$z^\iota = z^\iota([a, b], \mu) = \ell_{2^{\iota-1}a}(\mu) = \ell_{2^{\iota-1}b}(\mu) \in \partial U_\iota$$

called **dynamic root points**, indexed by  $\iota \in \mathbb{Z}/p$ .



The  $a/2$  and  $b/2$  rays land on *opposite* sides of  $U_0$ :

$$\ell_{a/2}(\mu) = \pm z^0([a, b], \mu) \quad \text{and} \quad \ell_{b/2}(\mu) = \mp z^0([a, b], \mu).$$



More generally, these periodic points  $z^\nu([a, b], \mu) \in \partial K(\hat{f}_\mu)$  are defined, and vary holomorphically with  $\mu$ , for all  $\mu$  in the wake  $W_{[a,b]}$ , even when  $\mu \notin \hat{M}$ .

Given any infinite sequence of hyperbolic components  $H_k$  of period  $p_k > 1$ , any point in the nested intersection,

$$\omega \in \bigcap_k (H_1 \triangleright \cdots \triangleright H_k \triangleright \widehat{M}),$$

is said to be ***infinitely renormalizable***. Let  $[a_k, b_k]$  be the characteristic interval for the  $k$ -fold tuning product


$$H_k^{\otimes} = H_1 \triangleright \cdots \triangleright H_k.$$

*I will consider only the case where the nested intersection  $\bigcap [a_k, b_k]$  consists of a single angle  $\theta$ . (An equivalent condition would be that  $H_k \cap \mathbb{R} = \emptyset$  for infinitely many  $k$ .)*

Note that the points

$$\mathbf{z}_k(\omega) = z^0([a_k, b_k], \omega)$$

and their negatives cut the filled Julia set  $K(\widehat{f}_\omega)$  into countably many pieces.



*In this schematic diagram, externals rays are orange, equipotentials are blue, and the Julia set is black.*

Let  $X$  be the connected component of zero in the set

$$K(\widehat{f}_\omega) \setminus \{\pm \mathbf{z}_k(\omega)\}.$$

**Lemma 1.**  *$X$  is compact, connected, and cellular. Every limit point of  $\{\pm \mathbf{z}_k(\omega)\}$ , and every limit point of the  $\theta/2$  and  $(1 + \theta)/2$  rays, belongs to  $X$ . But every other ray is bounded away from  $X$ . If  $K(\widehat{f}_\omega)$  is locally connected then  $X = \{0\}$ .*

**Conjecture.** *Conversely, if  $X = \{0\}$ , then  $K(\widehat{f}_\omega)$  is locally connected.*

## Choosing the Angles.

Now assume that the  $H_k$  are satellite hyperbolic components  $H(n_k/p_k)$ . Again let  $H_k^{\otimes} = H_1 \triangleright H_2 \triangleright \cdots \triangleright H_k$ , with characteristic interval  $[a_k, b_k]$ , and let  $r_k$  be the root point of  $H_k^{\otimes}$ . Then the periodic point


$$\mathbf{z}_j(r_k) = z^0([a_j, b_j], r_k) \in \partial K(\widehat{f}_{r_k}).$$

is defined for all  $j \leq k$ . (This point has period  $p_1 \cdots p_{j-1}$ . It is parabolic for  $j = k$ , and repelling for  $j < k$ .)

**Lemma 2.** *We can choose the angles  $t_j = n_j/p_j \neq 0$  inductively so that these points  $\mathbf{z}_j(r_k)$  are within some specified neighborhood of  $\mathbf{z}_1(r_1)$  for all  $j \leq k$ .*

**Start Proof.** Suppose  $H_1, \dots, H_k$  have already been chosen. We must show that each  $\mathbf{z}_j(r_{k+1})$  with  $j \leq k+1$  depends continuously on the choice of  $r_{k+1}$ , and hence can be placed arbitrarily close to  $\mathbf{z}_j(r_k)$  by choosing  $r_{k+1}$  close to  $r_k$ . 16





Since  $r_1, \dots, r_k$  have been chosen,  $\mathbf{z}_k(r_k)$  is a well defined parabolic point of period  $p_1 \cdots p_{k-1}$  and multiplier  $e^{2\pi i n_k / p_k}$ .

For  $\mu$  in a small neighborhood of  $r_k$ , the orbit of  $\mathbf{z}_k(r_k)$  splits into an orbit of the same period  $p_1 \cdots p_{k-1}$  with multiplier  $\approx e^{2\pi i n_k / p_k}$ , and a nearby orbit of period  $p_1 \cdots p_k$  with multiplier  $\approx +1$ .

Take  $\mu = r_{k+1}$  to be a point at rational angle along the boundary of  $H_k^*$ . Then this new orbit will again be parabolic, and the orbit point  $\mathbf{z}_{k+1}(r_{k+1})$  will converge to  $\mathbf{z}_k(r_k)$  as  $r_{k+1}$  converges to  $r_k$ .

Since the points  $\mathbf{z}_j(r_{k+1})$  with  $j \leq k$  clearly vary continuously with  $r_{k+1}$ , this proves Lemma 2.  $\square$

## Proof of the Theorem.

Recall that:


- The sequence  $\{r_k\}$  of root points has  $\omega$  as limit point.
- The function  $\mu \mapsto \mathbf{z}_j(\mu)$  is continuous for  $\mu \in W_{[a_j, b_j]}$ .

Therefore, for each fixed  $j$ , the sequence of points  $\mathbf{z}_j(r_k)$  has  $\mathbf{z}_j(\omega)$  as a limit point.


By Lemma 2, we can choose the  $H_k$  so that the  $\mathbf{z}_j(r_k)$  are uniformly bounded away from 0.

Hence the points  $\mathbf{z}_j(\omega)$  are also bounded away from 0.

Therefore, by Lemma 1,  $K(\hat{f}_\omega)$  cannot be locally connected.  $\square$




If  $t_k > 0$  for all angles  $t_k$ , then the  $\theta/2$  and  $(\theta + 1)/2$  rays spiral around each other without landing, in a “paper clip” pattern as sketched above, with the Julia set spiraling between them.



Here is a schematic picture close to the  $(\theta/2)$ -ray, which has been straightened out. All the points  $z_j(\omega)$  are assumed to lie in the region  $Z$ , while their negatives lie in  $-Z$ .

## The $\sin(1/x)$ model.



On the hand, if the signs of the  $t_k$  alternate, then the Julia set (indicated here in black) contains a  $\sin(1/x)$ -like curve. Compare Sørensen.

## How rapidly must $t_k \rightarrow 0$ ?

What rate of convergence is needed to guarantee that the  $\mathbf{z}_k(\omega)$  do **not** converge to zero? Here is a wild guess.

Perhaps there are order of magnitude estimates something like the following

$$\log \frac{\mathbf{z}_k(\omega)}{\mathbf{z}_{k+1}(\omega)} \approx \log \frac{\mathbf{z}_k(r_k)}{\mathbf{z}_{k+1}(r_{k+1})} \approx t_{k+1}^{1/p_k}$$

so that  $\{\mathbf{z}_k(\omega)\}$  converges to zero if and only if

$$\sum_k t_{k+1}^{1/p_k} = \infty. \quad (??)$$

For example, if  $t_k = 1/p_k$  with  $p_{k+1} = (k+1)^{p_k}$ , then

$$p_1 = 1, \quad p_2 = 2, \quad p_3 = 9, \quad p_4 = 262144, \quad p_5 \approx 1.2 \times 10^{27}, \quad \dots$$

tending rapidly to infinity. Yet  $t_{k+1}^{1/p_k} = 1/(k+1)$  with sum  $+\infty$ .

Conjecturally, this  $\{p_k\}$  does not increase fast enough! 21


## To conclude: Four Pictures

It is probably impossible to make any real picture of one of these non locally-connected Julia sets. However, we may get some intuitive idea by looking at relatively modest iterated satellite tunings.


In the first two pictures, the separating periodic points  $\mathbf{z}_1$  and  $\mathbf{z}_2$  are circled. The rays of angle  $a_1/2 = 1/14$  and  $b_1/2 = 1/7$  are shown, but those of angle  $a_2/2 \approx b_2/2$  are too close to distinguish from  $1/14$  respectively  $1/7$ .

In the last two pictures,  $\mathbf{z}_1$ ,  $\mathbf{z}_2$  and  $\mathbf{z}_3$  are defined and circled. (As the angles  $t_2, t_3$  tend to zero, these circled points would converge towards each other.) In these cases, the rays of angle  $a_1/2 = 1/14 < a_2/2 < b_2/2 < b_1/2 = 1/7$  can be distinguished.

(Assuming only that  $p_k \geq 3$  for all  $k$ , it follows that the differences  $b_k - a_k \approx 2^{-p_1 \cdots p_k}$  tend faster than exponentially to zero as  $k \rightarrow \infty$ .)




$H(1/3) \triangleright H(1/20) \triangleright 0$




$$H(1/3) \triangleright H(-1/20) \triangleright 0$$














$H(1/3) \triangleright H(1/7) \triangleright H(1/13) \triangleright 0$



$$H(1/3) \triangleright H(-1/7) \triangleright H(1/13) \triangleright 0$$

## References

-  A. Douady, Algorithms for computing angles in the Mandelbrot set, pp. 155–168 of “Chaotic Dynamics and Fractals”, ed. Barnsley & Demko, Acad. Press, 1986
-  A. Douady, Chirurgie sur les applications holomorphes, Proc. Int. Cong. Math. Berkeley, A.M.S. 1987, 724–738.
-  A. Douady and J. H. Hubbard, On the dynamics of polynomial-like mappings, Ann. Sci. Ec. Norm. Sup. (Paris) **18** (1985) 287–343.
-  C. McMullen, “Complex Dynamics and Renormalization,” Princeton U. Press, 1994.
-  J. Milnor, Self-similarity and hairiness in the Mandelbrot set, in “Computers in Geometry and Topology,” (M.C. Tangora, ed.), Marcel Dekker, New York, Basel, 1989, pp 211–257.

-  J. Milnor, Local connectivity of Julia sets: expository lectures, pp. 67–116 of “The Mandelbrot set, Theme and Variations” edit: Tan Lei, LMS Lecture Note Series **274**, Cambridge U. Press, 2000.
-  J. Milnor, Periodic orbits, external rays and the Mandelbrot set: An expository account, *Asterisque* **261** (2000) “Geometrie Complexe et Systemes Dynamiques”, pp. 277–333.
-  J. Milnor, “Dynamics in One Complex Variable,” (3rd ed.) Princeton U. Press, 2006.
-  D. E. K. Sørensen, Infinitely renormalizable quadratic polynomials, with non-locally connected Julia set, *J. Geom. Anal.* **10** (2000), 169–206