

# Math53: Ordinary Differential Equations Winter 2004

## Homework Assignment 3

**Problem Set 3 is due by 2:15p.m. on Monday, 2/9, in 380Y**

### Problem Set 3:

5.1: 14,26; 5.2:24,32,43; 5.3: 2,30; 5.4: 18,36; 5.5: 20; 5.6: 1,8; 5.7: 14,22,28  
PS3-Problems 16,17 (see next page)

*Note 1:* While there are two non-textbook problems this time, they should involve less writing than the non-textbook problems on the first two problem sets did. The first problem is meant to complement Section 5.6 in the book.

*Note 2:* Since this problem set is due on a Monday, I will have office hours 4-6p.m. on Sunday, 2/8.

### Daily Assignments:

<i>Date</i>	<i>Read</i>	<i>Exercises</i>
1/30 F	5.1, 5.2	5.1:14,26; 5.2:24,32,43
2/2 M	5.3	5.3:2,30
2/3 T	5.4,5.5	5.4:18,36; 5.5:20
2/4 W	5.6,5.7	5.6:1,8; PS3-Problem 16
2/5 R		5.7:14,22,28; PS3-Problem 17
2/6 F	5.8	

*Note:* You do not need to memorize Table 1 on p250. It will be put on the second midterm and on the final exam if it is needed for any problem. However, you should know how to use it and how to compute the Laplace transforms that appear in this table yourself.

### PS3-Problem 16

A smooth function  $f$  defined on the entire real line is called *compactly supported* if there exist  $a$  and  $b$  such that  $f(t) = 0$  for all  $t < a$  and for all  $t > b$ . In other words, compactly supported functions are nonzero only in the "middle" of the real line. The set of all such functions is denoted by  $C_c^\infty(\mathbb{R})$ . A *distribution* on  $\mathbb{R}$  is any linear operator  $T : C_c^\infty(\mathbb{R}) \rightarrow \mathbb{R}$ . In other words, for every compactly supported function  $f$ ,  $Tf$  is a real number, and

$$T(\alpha f_1 + \beta f_2) = \alpha T(f_1) + \beta T(f_2)$$

for all real numbers  $\alpha$  and  $\beta$  and compactly supported functions  $f_1$  and  $f_2$ . For example, the three operators given by

$$T(f) = 0, \quad T(f) = f'(1) - f(2), \quad T_\delta(f) = f(0),$$

are distributions.

(a) If  $g$  is a piecewise continuous function on the real line, show that the operators given by

$$T_g f = \int_{-\infty}^{\infty} g(t)f(t) dt, \quad f \in C_c^\infty(\mathbb{R}), \quad \text{and} \quad T'_g f = - \int_{-\infty}^{\infty} g(t)f'(t) dt \quad f \in C_c^\infty(\mathbb{R}),$$

are well-defined distributions.

(b) Use integration by parts to show that if  $g$  is a continuous function with a piecewise continuous first derivative and  $H$  is the Heaviside function, then

$$T'_g f = T_{g'} f \quad \text{and} \quad T'_H f = T_\delta f \quad \text{for all } f \in C_c^\infty(\mathbb{R}).$$

If  $g_1 \neq g_2$  are piecewise continuous functions, then  $T_{g_1} \neq T_{g_2}$  as operators, i.e. the numbers  $T_{g_1}f$  and  $T_{g_2}f$  are not the same for at least one  $f \in C_c^\infty(\mathbb{R})$ . Thus, every such function  $g$  can be thought of as the distribution  $T_g$ . Furthermore, if  $g$  is smooth,  $T'_g = T_{g'}$  by (b). However, not every distribution is of the form  $T_g$ , for some piecewise continuous function  $g$ . For example,  $T_\delta$  is not of this form. In this sense, distributions are *generalized functions*, and in the sense of distributions  $H' = \delta$ , i.e.  $T'_H = T_\delta$ . The limit in Definition 6.2 on p282, means that

$$\lim_{\epsilon \rightarrow 0} T_{\delta_\epsilon} f = T_\delta f \quad \text{for all } f \in C_c^\infty(\mathbb{R}).$$

### PS3-Problem 17

Recall from PS2-Problem 1, that if  $p$  and  $q$  are constants,

$$(e^{(\lambda_2 - \lambda_1)t} (e^{-\lambda_2 t} y)')' = e^{-\lambda_1 t} (y'' + py' + qy), \quad (1)$$

where  $\lambda_1$  and  $\lambda_2$  are the two roots of the characteristic polynomial  $\lambda^2 + p\lambda + q = 0$  associated to the linear homogeneous second-order ODE  $y'' + py' + qy = 0$ .

(a) Use (1) to find a solution  $y_p = y_p(t)$  to the ODE

$$y'' + py' + qy = f, \quad f = f(t),$$

in the form  $y_p = G * f$  for some function  $G = G(t)$ . In particular, give an explicit expression for  $G$ . Use only (1) to find  $y_p$ , and not Section 5.7 in the book. You will need to consider two cases in your computation:  $\lambda_1 = \lambda_2$  and  $\lambda_1 \neq \lambda_2$ . In the latter case, there will be two subcases. In one of these subcases, the expression you obtain for  $G = G(t)$  will involve complex numbers, but simplifies to a real expression.

(b) Compare your expression for  $y_p$  with that for  $y_s$  in Theorem 7.16, on p293. In particular, use the inverse Laplace transform and Theorem 6.10, on p285, to give an explicit expression for the function  $e = e(t)$  that appears in the definition of  $y_s$  in each of the three cases.