

# MAT 550 – Test 2 – Spring 2009

You may use any results proved in the book.  $\hat{\cdot}$  denotes the Fourier transform.

1. (a) Let  $\{\psi^t\}$  be a standard approximate identity on  $\mathbf{R}^n$ . Show that, if  $f \in L^1(\mathbf{R}^n)$ , then  $f * \psi^t \rightarrow f$  in  $L^1$  as  $t \rightarrow 0^+$ .

(Hint: you can use the homework Exercises 11.4.8 and 11.4.9 (attached).)

- (b) Say  $f \in L^1(\mathbf{R}^n)$  and  $\int f\varphi = 0$  for every  $\varphi \in \mathcal{S}(\mathbf{R}^n)$ ; show that  $f = 0$ .

(Hint: use (a).)

- (c) According to a proposition in the book, if  $F, G \in L^1(\mathbf{R}^n)$ , then  $\int \hat{F}\bar{G} = \int F\bar{\hat{G}}$ . Use this and (b) to show:

If  $f \in L^1(\mathbf{R}^n)$ , and also  $\hat{f} \in L^1(\mathbf{R}^n)$ , then  $f = (\hat{f})^\vee$ .

- (d) Conclude that  $\hat{\cdot}$  is injective on  $L^1(\mathbf{R}^n)$ .

**Solution** (a) A standard approximate identity has the form  $\psi_t(x) = t^{-n}\psi(x/t)$  for some  $\psi \in C_c^\infty$  with  $\psi \geq 0$  and  $\int \psi = 1$ . Choose  $R$  such that the support of  $\psi$  is contained in  $\bar{B}_R$ , where  $B_R$  is the open ball of radius  $R$  centered at 0. Then for  $0 < t < 1$ ,  $\text{supp}\psi_t \subseteq \bar{B}_{tR} \subseteq \bar{B}_R$ . Let  $K_2 = \bar{B}_R$ .

Now say first  $f \in C_c$ , and say  $\text{supp}f = K_1$ . Then for  $0 < t < 1$ ,  $\text{supp}(f*\psi_t) \subseteq K_1 + K_2 = K$ , say. Also  $f * \psi_t \rightarrow f$  uniformly (as  $t \rightarrow 0^+$ ). Now  $\|f * \psi_t - f\|_1 \leq m(K)\|f * \psi_t - f\|_\infty$ , so in this case, we do have  $f * \psi_t \rightarrow f$  in  $L^1$ .

But  $C_c$  is dense in  $L^1$ , so by the fundamental maximal principle, we need only prove the following. Define  $T_t : L^1 \rightarrow L^1$  by  $T_t f = f * \psi_t$ ; then  $\|T_t\| \leq 1$  for all  $t$ . But this is clear, since  $\|T_t f\|_1 \leq \|f\|_1 \|\psi_t\|_1$ , and  $\int f\psi_t = \int f\psi = 1$  for all  $t$ .

- (b) In the notation of (a), it is clearly enough to show that  $f * \psi_t \equiv 0$  for all  $t$ . But  $f * \psi_t(x) = \int f(y)\psi_t(x-y)dy$ , and for any fixed  $x$ , the function  $y \rightarrow \psi_t(x-y)$  is clearly in  $\mathcal{S}(\mathbf{R}^n)$ . By the hypotheses of (b), this is zero.

- (c) Actually it's a little easier to use the equivalent fact that if  $F, G \in L^1(\mathbf{R}^n)$ , then  $\int \hat{F}G = \int F\hat{G}$ .

To show that  $f - (\hat{f})^\vee = 0$ , it is enough to show, by (b), that for all  $\varphi \in \mathcal{S}$ ,  $\int (\hat{f})^\vee \phi = \int f\phi$ . But

$$\int (\hat{f})^\vee \phi = \int \hat{f}\hat{\phi} = \int f(\hat{\phi})^\vee = \int f\phi$$

as claimed. (We used the inversion theorem for  $\mathcal{S}$ .)

- (d) Say  $\hat{f} = 0$ . Note that  $0 \in L^1$ . Thus, by (c),  $f = (\hat{f})^\vee = \check{0} = 0$ .

2. Suppose  $a, b, c$  are real numbers, and that  $b^2 - ac > 0$ . Find, with proof, the general solution of the partial differential equation

$$au_{xx} + 2bu_{xy} + cu_{yy} = 0.$$

Here  $u(x, y)$  is to be a  $C^2$  function on  $\mathbf{R} \times \mathbf{R}$ , and the subscripts denote differentiation. (Hint: what special case have we done, and how did we do it?)

**Solution** If  $a = c = 0$ , the general solution is  $u = h_1(x) + h_2(y)$  for  $C^2$  functions  $h_1, h_2$ . Otherwise assume without loss that  $a \neq 0$ . A special case is the wave equation, in which  $b = 0, a = 1, c = -1$ , and we proceed similarly. Writing the quadratic  $aX^2 + 2bX + c = a(X - r_1)(X - r_2)$ , where the real roots  $r_1, r_2$  are given by the quadratic equation formula, we find

$$a(D_x - r_1 D_y)(D_x - r_2 D_y)u = 0.$$

We change coordinates to  $(p, q)$  so that  $\partial/\partial p = D_x - r_1 D_y, \partial/\partial q = D_x - r_2 D_y$ ; by the chain rule we want  $x = p + q, y = -r_1 p - r_2 q$ . Inverting the matrix of this linear transformation we find  $p = -D(r_2 x + y), q = D(r_1 x + y)$ , where  $D = (r_1 - r_2)^{-1}$ . The equation is now  $\partial^2 u / \partial p \partial q = 0$ , whose general solution is  $u = h_1(p) + h_2(q) = H_1(r_2 x + y) + H_2(r_1 x + y)$  for  $C^2$  functions  $H_1, H_2$ .

When  $b^2 - ac > 0, ax^2 + 2bxy + cy^2$  is the equation of a hyperbola, so the PDE is called *hyperbolic*. Similarly it is called *elliptic* if  $b^2 - ac < 0$ .

3. Let  $\mathbf{R}_+^2$  be the open upper half-plane,  $\{(x, y) \in \mathbf{R}^2 : y > 0\}$ , and let  $\overline{\mathbf{R}_+^2}$  be its closure. Find a solution of the Dirichlet problem for  $\mathbf{R}_+^2$ , in the following sense: Let  $f$  be a bounded continuous function on  $\mathbf{R}$ . Find, with proof, a function  $u(x, y)$  which is continuous on  $\overline{\mathbf{R}_+^2}$ , and which satisfies

$$\Delta u = 0 \text{ on } \mathbf{R}_+^2, \text{ and}$$

$$u(x, 0) = f(x) \text{ for all } x \in \mathbf{R}.$$

Express your solution in the form  $u(x, y) = (f * \varphi_y)(x)$  for a completely explicit function  $\varphi_y$ . (Hint: first conjecture a solution, by using the Fourier transform in the  $x$  variable.) **Do this for homework.**