

# H.W. # 8


2)  $\int_0^{\pi/2} \cos^5 x dx = \int_0^1 (1-u^2)(1-u^2) du$ ;  $u = \sin x$   $\rightarrow \begin{cases} x = \frac{\pi}{2} \rightarrow u = 1 \\ x = 0 \rightarrow u = 0 \end{cases}$

$\int_0^1 (1-2u^2+u^4) du = u - \frac{2u^3}{3} + \frac{u^5}{5} \Big|_0^1 = 1 - \frac{10}{15} + \frac{3}{15} = \frac{8}{15}$

8)  $\int_0^{\pi/4} \tan^2 x \sec^4 x dx = \int_0^1 u^2(u^2+1) du$ ;  $u = \tan x$   $\rightarrow \begin{cases} x = 0 \rightarrow u = 0 \\ x = \frac{\pi}{4} \rightarrow u = 1 \end{cases}$

$= \int_0^1 (u^4 + u^2) du = \frac{u^5}{5} + \frac{u^3}{3} \Big|_0^1 = \frac{1}{5} + \frac{1}{3} = \frac{8}{15}$

10) Using  $x = \sec \theta$   
 $dx = \sec \theta \tan \theta d\theta \rightarrow \int \frac{\sqrt{x^2-1}}{x^4} dx = \int \frac{\sqrt{\sec^2 \theta - 1}}{\sec^3 \theta} \tan \theta d\theta = \int \frac{\tan^2 \theta}{\sec^3 \theta} d\theta = \int \frac{\sin^2 \theta}{\cos^3 \theta} \cdot \cos^3 \theta d\theta$   
 $= \int \sin^2 \theta \cos \theta d\theta = \int u^2 du = \frac{u^3}{3} + C = \frac{\sin^3 \theta}{3} + C$

Using  $x = \sec \theta$    $\rightarrow \sin \theta = \frac{\sqrt{x^2-1}}{x}$

1)  $\int_0^1 \frac{x-1}{x^2+3x+2} dx \rightarrow \left( \frac{x-1}{x^2+3x+2} = \frac{A}{x+2} + \frac{B}{x+1} \right) (x+2)(x+1)$

$\Rightarrow x-1 = A(x+1) + B(x+2)$

$x-1 = x(A+B) + (A+2B)$

Using  $A+B=1$  &  $A+2B=-1 \rightarrow -A=-3 \rightarrow A=3$  and  $B=-2$

$\int_0^1 \frac{3}{x+2} dx - \int_0^1 \frac{2}{x+1} dx = 3 \ln|x+2| \Big|_0^1 - 2 \ln|x+1| \Big|_0^1 = 3(\ln 3 - \ln 2) - 2 \ln 2 = 3 \ln 3 - 5 \ln 2$

$$(8.) \int_0^1 \frac{x^3 - 4x - 10}{x^2 - x - 6} dx$$

$$\begin{array}{r}
 x^2 - x - 6 \overline{) x^3 + 0x^2 - 4x - 10} \\
 \underline{-(x^3 - x^2 - 6x)} \phantom{-10} \\
 0 \phantom{x^2} + 2x - 10 \\
 \underline{-(x^2 - x - 6)} \\
 0 \phantom{x^2} + 3x - 4
 \end{array}
 + \frac{3x-4}{x^2-x-6}$$

$$\int_0^1 (x+1) dx + \int_0^1 \frac{3x-4}{x^2-x-6} dx$$

$$\frac{x^2 + x \Big|_0^1}{2} = \frac{3}{2}$$

$$\frac{3x-4}{x^2-x-6} = \frac{A}{x-3} + \frac{B}{x+2}$$

$$3x-4 = A(x+2) + B(x-3)$$

$$3x-4 = x(A+B) + (2A-3B) \rightarrow \begin{cases} A+B=3 \\ 2A-3B=-4 \end{cases}$$

Plugging in, we get  $2A-3(3-A)=-4 \rightarrow 5A=5 \rightarrow A=1$  and  $B=2$

$$\int_0^1 \frac{3x-4}{x^2-x-6} dx = \int_0^1 \frac{1}{x-3} dx + \int_0^1 \frac{2}{x+2} dx = \ln|x-3| \Big|_0^1 + 2 \ln|x+2| \Big|_0^1 = \ln 2 - \ln 3 + 2(\ln 3 - \ln 2) = \ln 3 - \ln 2$$

$$\int_0^1 \frac{x^3 - 4x - 10}{x^2 - x - 6} dx = \frac{3}{2} + \ln 3 - \ln 2$$

or

$$= \frac{3}{2} + \ln\left(\frac{3}{2}\right)$$

5.9 # 2, 18

2-) From the concavity of  $y=f(x)$ , we see that  $\begin{cases} L_n \text{ left end approx.} \rightarrow \text{overestimate } (0.9540) \\ R_n \text{ right end approx.} \rightarrow \text{underestimate } (0.7811) \end{cases}$

We can easily distinguish  $T_n$  from  $M_n$  by remembering (pg. 413) that:  
trapezoidal approx:  $T_n = \frac{R_n + L_n}{2} = \frac{0.9540 + 0.7811}{2} \approx 0.8675$

Since  $R_n < M_n < L_n$  and  $R_n < T_n < L_n$ , we conclude that:

$$T_n = 0.8675 \ \& \ M_n = 0.8632$$

(b) It should be clear that  $M_n, T_n$  are much closer to  $\int_0^2 f(x) dx$  compared with  $L_n, R_n$ .

- If  $f(x)$  is concave up from  $x=0$  to  $x=2$ , then:  $M_n < \int_0^2 f(x) dx < T_n$

- If  $f(x)$  is concave down from  $x=0$  to  $x=2$ , then  $T_n < \int_0^2 f(x) dx < M_n$

Since our graph ( $y=f(x)$ ) is concave up:

$$0.8632 < \int_0^2 f(x) dx < 0.8675$$

18-) (a) Using eq. on pg. 413:

$$T_8 = \frac{1}{16} [f(0) + 2f(\frac{1}{8}) + 2f(\frac{2}{8}) + 2f(\frac{3}{8}) + 2f(\frac{4}{8}) + 2f(\frac{5}{8}) + 2f(\frac{6}{8}) + 2f(\frac{7}{8}) + f(1)] \quad \begin{matrix} a=0 \rightarrow x_i = i\Delta x \\ \Delta x = \frac{1}{8} \end{matrix}$$

$$= \frac{1}{16} [1 + 2\cos(\frac{1}{64}) + 2\cos(\frac{1}{16}) + \dots + \cos 1]$$

$$T_8 \approx 0.9023$$

Using midpoint rule ( $\bar{x}_i = \frac{1}{2}(x_{i-1} + x_i)$ ):

$$M_8 = \frac{1}{8} [f(\frac{1}{16}) + f(\frac{3}{16}) + f(\frac{5}{16}) + f(\frac{7}{16}) + f(\frac{9}{16}) + \dots + f(\frac{15}{16})]$$

$$M_8 \approx 0.9056$$


(b) We want to apply eq. 3 for errors, so:

Using chain rule for  $f(x) = \cos(x^2) \rightarrow f'(x) = -2x(\sin(x^2)) \xrightarrow{\text{product rule}} f''(x) = \frac{-2\sin(x^2) + 4x^2\cos(x^2)}{-2\sin(x^2) - 4x^2\cos(x^2)}$

$$|f''(x)| = 2(\sin x^2 + 2x^2\cos x^2)$$

Using  $a=0, b=1 \rightarrow$  For  $0 \leq x \leq 1 \rightarrow \boxed{x^2 \leq 1} \rightarrow \boxed{\sin x^2 \leq 1 \ \& \ \cos x^2 \leq 1}$

So we now have:  $|f''(x)| \leq 2(1+2 \cdot 1) = 6 \rightarrow K=6$  (from eq. 3)



From trapezoid rule:  $(a=0, b=1, n=8)$

$$|E_T| \leq \frac{K(b-a)^3}{12n^2} = \frac{6(1-0)^3}{12(8)^2} = \frac{1}{128} \rightarrow |E_T| \leq 0.00781$$

From midpt. rule:

$$|E_M| \leq \frac{K(b-a)^3}{24n^2} = \frac{1}{2} \left( \frac{1}{128} \right) = \frac{1}{256} \rightarrow |E_T| \leq 0.00391$$

c-) From trap. rule:  $\frac{6(1)^3}{12n^2} < 0.00001 \rightarrow n^2 > \frac{1}{0.00002} \rightarrow n > 223.61 \rightarrow n \geq 224$  <sup>optional</sup>

From midpt. rule:  $\frac{6(1)^3}{24n^2} < 0.00001 \rightarrow n^2 > \frac{1}{4}(10^5) \rightarrow n > 158.11 \rightarrow n \geq 159$  <sup>optional</sup>