

5. Suppose a lottery ticket consists of six numbers from 1 to 44. The six winning numbers are chosen randomly. Which of the following tickets has the best chance of winning?
- (A) 11 12 13 14 15 16
 - (B) 09 13 22 25 31 43
 - (C) 17 21 33 34 38 41

Did you recognize that each of the lottery tickets in Question #5 has the same chance of winning? Most people would avoid buying a ticket like (A) because it is not "random" enough – they realize that the odds against the ticket winning are very great. In fact, these people are right. What they might not realize is that the odds are *equally* great against either (B) or (C) winning – the odds against *any* ticket winning are 7,059,051 to 1. When it comes to randomness, intuition can be a dangerous thing to trust.

How You Can Get Random

Let us see how you might construct a random sequence of 26 B's and 26 R's as you were asked to do in Question #3.

6. Separate a deck of cards into a pile of black cards and a pile of red cards. Then flip a fair coin repeatedly to form your sequence of cards; if the coin lands on heads, choose a black card, and if the coin lands on tails, choose a red card. When one of the piles runs out of cards, finish the sequence with the remaining cards in the other pile. You have just generated a random sequence of black and red cards!
7. Perform the same experiment as in question #6, but this time enter the command **randInt(1,2)** on your calculator and press **ENTER** repeatedly. (To find **randInt**(press **MATH** **◀** **5**). If the output is 1, choose a black card, and if the output is 2, choose a red card. When one of the piles runs out of cards, finish the sequence with the remaining cards in the other pile. This sequence is also a random sequence.
8. If you have a TI-83/84 Plus calculator with the **Prob Sim** App, you can watch the calculator draw cards for you. (The **Prob Sim** App can be downloaded for free at http://education.ti.com/educationportal/sites/US/productDetail/us_prob_sim_83_84.html.) Choose the **Draw Cards** option from the menu, select **SET**, and answer **No** to **Replace**. Press the **OK** button. You can then press **ENTER** repeatedly to draw cards in a random order. (Remember, clubs and spades are black cards, and hearts and diamonds are red cards.)

The randomness in Question #6 relies on heads and tails being equally probable every time you flip the coin, an assumption that might not be valid. The randomness in Questions #7 and #8 relies on your calculator's ability to generate random numbers, which it can do only because it has a *pseudo-random number generator* programmed into its software. (For more about pseudo-random number generators, see the Extensions page of this activity.)

The goal of this activity is to give your students a short and simple snapshot into a very extensive math topic. TI and NCTM encourage you and your students to learn more about this topic using the extensions provided below and through your own independent research.

Extensions

More Probability Simulations

- A fair 6-sided die (or number cube) has an equal probability of showing any of the numbers 1 through 6. Use your calculator to generate a random sequence of 50 rolls of a fair die. How many times did you roll the same number three times in a row?
- According to www.AABB.org, 45% of blood donors have type O blood, 40% have type A blood, 11% have type B blood, and 4% have type AB blood. Use your calculator to generate a random sequence of 20 donors classified by blood type. [Hint: Generate numbers between 00 and 99. Let 00 through 44 represent type O, since those are 45% of the possible numbers. Assign the remaining numbers to blood types according to their percentages.]

Suppose a mobile blood unit processes 40 donors in a typical day. Is it unusual for the unit to finish a day with no units of type AB blood? Use your calculator to perform your simulation ten times, each time for 40 donors, and see how many include no AB donors.

Random and Pseudo-Random Numbers

The ability of computers to simulate random processes is useful in many modern applications, ranging from the obvious (e.g., computer games and cryptography) to the less obvious (e.g., weather forecasting and fighting crime). In some applications, particularly cryptography (code design), pseudo-random numbers are dangerously predictable, so "true" random number generators are preferred. These generators derive numerical data from naturally-occurring sources of entropy, two of the best-known examples of which are atmospheric noise and radioactive decay. Other sources that have been used include lava lamps.

- One of the most famous books in applied mathematics is *A Million Random Digits with 100,000 Normal Deviates*, published in 1955 by the RAND Corporation. As the title suggests, it is a book consisting almost entirely of numbers. An intriguing history of the book, along with a sampling of the statistical tests to which it was subjected to prove its true randomness, can be found at www.rand.org/publications/classics/randomdigits.
- An article by Ivars Peterson about how SGI used lava lamps to generate random numbers can be found at www.maa.org/mathland/mathtrek_5_7_01.html.
- The Web site www.random.org has a nice explanation of pseudo-random numbers and true random numbers. It is also a source for true random numbers in various useful forms (they use atmospheric noise to generate random numbers).
- You can find out more about pseudo-random number generators in the activity **Creating "Random" Numbers** that accompanied the *NUMB3RS* episode "Bettor or Worse."