

Your name: _____

TA's name: _____

Problem #1: Find the derivative of each function.

$$a) f(x) = \frac{3-\sqrt{x}}{3+\sqrt{x}}$$

$$f'(x) = \frac{(3+\sqrt{x})\left(\frac{1}{2\sqrt{x}}\right) - (3-\sqrt{x})\left(\frac{1}{2\sqrt{x}}\right)}{(3+\sqrt{x})^2}$$

$$b) f(x) = (3x^2 + 9x - 4)(4x^3 + x^2 - x)$$

$$f'(x) = (3x^2 + 9x - 4)(12x^2 + 2x - 1) + (4x^3 + x^2 - x)(6x + 9)$$

Problem #2: Find the equation of the tangent line to $y = \sin(4x)$ at $x = \frac{\pi}{16}$.

$$y = \sin\left(\frac{\pi}{16}\right) = \sin\frac{\pi}{4} = \frac{\sqrt{2}}{2}$$

$$y - \frac{\sqrt{2}}{2} = m(x - \frac{\pi}{16})$$

$$\frac{dy}{dx} = \cos(4x)(4)$$

$$\text{at } x = \frac{\pi}{16} : \frac{dy}{dx} = \cos\left(4\left(\frac{\pi}{16}\right)\right)(4) = 4\left(\frac{\sqrt{2}}{2}\right) = 2\sqrt{2}$$

$$y - \frac{\sqrt{2}}{2} = 2\sqrt{2}(x - \frac{\pi}{16})$$

Problem #3. Find all x -values of

$$f(x) = x^3 - 6x^2 - 36x + 9$$

for which either $f'(x) = 0$ or $f'(x)$ is not defined.

$$\begin{aligned} f'(x) &= 3x^2 - 12x - 36 = 0 \\ x^2 - 4x - 12 &= 0 \\ (x-6)(x+2) &= 0 \\ x = 6 \quad x = -2 \end{aligned}$$

Problem #4: Find $\frac{dy}{dx}$ if $x^3 - 5xy^2 + y^3 = 1$.

$$3x^2 - [5x(2y \frac{dy}{dx}) + 5y^2] + 3y^2 \frac{dy}{dx} = 0$$

$$3x^2 - 10xy \frac{dy}{dx} - 5y^2 + 3y^2 \frac{dy}{dx} = 0$$

$$3x^2 - 5y^2 = 10xy \frac{dy}{dx} - 3y^2 \frac{dy}{dx}$$

$$3x - 5y^2 = \frac{dy}{dx} (10xy - 3y^2)$$

$$\frac{3x - 5y^2}{10xy - 3y^2} = \frac{dy}{dx}$$

Problem #5: Find the equation of the tangent line to $\ln(2x^2 - y^2) = 0$ at $(1,1)$.

$$y-1 = m(x-1)$$

$$\frac{4x-2y \frac{dy}{dx}}{2x^2-y^2} = 0$$

$$\frac{4-2 \frac{dy}{dx}}{2-1} = 0$$

$$4-2 \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} = 2$$

$$y-1 = 2(x-1)$$

Problem #6: Find $\frac{dy}{dx}$ if:

(a) $y = \tan^{-1}(2x)$

$$\frac{dy}{dx} = \frac{1}{1+(2x)^2} \cdot 2 = \frac{2}{1+4x^2}$$

b) $f(x) = \sin^3\left(\frac{2-5x}{x^2}\right)$

$$f'(x) = 3 \sin^2\left(\frac{2-5x}{x^2}\right) \left[\frac{x^2(-5)}{x^4} - \frac{(2-5x)(2x)}{x^4} \right]$$

Problem #7: Find the points (x, y) where the line tangent to

$$y = x^3 - 6x^2 - 30x + 4$$
 is parallel to $15x + y = 10$.

$$\begin{aligned}15x + y &= 10 \\y &= -15x + 10 \\ \text{slope} &= -15\end{aligned}$$

$$\frac{dy}{dx} = 3x^2 - 12x - 30 \leftarrow \text{parallel so slopes are equal}$$

$$3x^2 - 12x - 30 = -15$$

$$3x^2 - 12x - 15 = 0$$

$$x^2 - 4x - 5 = 0$$

$$(x-5)(x+1) = 0$$

$$x = 5, x = -1$$

$$y = 5^3 - 6(5^2) - 30(5) + 4$$

$$= 125 - 150 - 150 + 4$$

$$= -171$$

$$(5, -171)$$

$$y = (-1)^3 - 6(-1)^2 - 30(-1) + 4$$

$$= -1 - 6 + 30 + 4$$

$$= 27$$

$$(-1, 27)$$

Problem #8. Find all values of x where $y = x^2e^x$ has an absolute maximum or minimum on the interval $[-3, 1]$.

This problem requires material we have not yet covered.

Problems like this will not be on the midterm.