

MAT 126

Solutions to Midterm 2 (acoustic)

1. Determine these EASY antiderivatives. You should be able to do these **very well**. In these problems, no justification is needed. Remember the '+C'.

6 pts

(a) $\int \frac{3}{x} dx$

Solution:

$$\int \frac{3}{x} dx = 3 \ln |x| + C$$

6 pts

(b) $\int 4 \sin(x) dx$

Solution:

$$\int 4 \sin(x) dx = -4 \cos(x) + C$$

6 pts

(c) $\int e^{3x} dx$

Solution:

$$\int e^{3x} dx = \frac{1}{3} e^{3x} + C$$

6 pts

(d) $\int \frac{3}{t^2 + 1} dt$

Solution:

$$\int \frac{3}{t^2 + 1} dt = 3 \arctan(t) + C$$

6 pts

(e) $\int \frac{1}{\sqrt{1-u^2}} du$

Solution:

$$\int \frac{1}{\sqrt{1-u^2}} du = \arcsin(u) + C$$

2. In this question we tell you which method we suggest you use. Use the back of the previous page if you need more space.

15 pts

(a) Suggested method: substitution $\int \frac{z}{1+z^2} dz$

Solution: Make the substitution $u = 1 + z^2$, so that $du = 2dz$, or $\frac{1}{2}du = dz$. Then

$$\int \frac{z}{1+z^2} dz = \frac{1}{2} \int \frac{1}{u} du = \frac{1}{2} \ln |u| + C = \boxed{\frac{1}{2} \ln |1+z^2| + C}$$

Note that since $1 + z^2 > 0$ for all z , the absolute value is not necessary; the answer $\frac{\ln(1+z^2)}{2} + C$ is fine, too.

15 pts

(b) Suggested method: substitution $\int \frac{e^{\sqrt{y+2}}}{\sqrt{y+2}} dy$

Solution: Make the substitution $u = \sqrt{y+2}$. Then

$$du = \frac{1}{2\sqrt{y+2}} dy \quad \text{or} \quad 2du = \frac{dy}{\sqrt{y+2}}$$

Thus,

$$\int \frac{e^{\sqrt{y+2}}}{\sqrt{y+2}} dy = 2 \int e^u du = 2e^u + C = \boxed{2e^{\sqrt{y+2}} + C}$$

15 pts

(c) Suggested method: substitution $\int \frac{\ln(x)}{x} dx$

Solution: Here, we let $u = \ln(x)$ and so $du = \frac{dx}{x}$. This means we have

$$\int \frac{\ln(x)}{x} dx = \int u du = \frac{u^2}{2} + C = \boxed{\frac{(\ln(x))^2}{2} + C}$$

3. In this question we tell you which method we suggest you use. Use the back of the previous page if you need more space.

15 pts

- (a) Suggested method: integration by parts $\int x^5 \ln(x) dx$

Solution: Take $u = \ln(x)$ and $dv = x^5 dx$. Then $du = \frac{1}{x} dx$ and $v = \frac{x^6}{6}$. So:

$$\int x^5 \ln(x) dx = \frac{x^6 \ln(x)}{6} - \frac{1}{6} \int x^6 \cdot \frac{1}{x} dx = \frac{x^6 \ln(x)}{6} - \frac{1}{6} \int x^5 dx = \boxed{\frac{x^6 \ln(x)}{6} - \frac{x^6}{36} + C}$$

15 pts

- (b) Suggested method: integration by parts $\int x e^{2x} dx$

Solution: Take $u = x$ and $dv = e^{2x} dx$. Then $du = dx$ and $v = \frac{1}{2} e^{2x}$, and so we have

$$\int x e^{2x} dx = \frac{x}{2} e^{2x} - \frac{1}{2} \int e^{2x} dx = \boxed{\frac{x e^{2x}}{2} - \frac{e^{2x}}{4} + C}$$

15 pts

- (c) Suggested method: integration by parts $\int \sin(x) e^{2x} dx$

Solution: Take $u = e^{2x}$ and $dv = \sin(x) dx$. Then $du = 2e^{2x} dx$ and $v = -\cos(x)$. So we have

$$\int \sin(x) e^{2x} dx = -\cos(x) e^{2x} + 2 \int \cos(x) e^{2x} dx$$

(we have a + before the integral because we were subtracting a negative). To do the second integral, we take $u = e^{2x}$ and $dv = \cos x dx$. Then $du = 2e^{2x} dx$ and $v = \sin x$. This gives us

$$\int \sin(x) e^{2x} dx = -\cos(x) e^{2x} + 2 \left(\sin(x) e^{2x} - 2 \int \sin(x) e^{2x} dx \right)$$

Multiplying out gives

$$\int \sin(x) e^{2x} dx = -\cos(x) e^{2x} + 2 \sin(x) e^{2x} - 4 \int \sin(x) e^{2x} dx$$

or, equivalently,

$$5 \int \sin(x) e^{2x} dx = -\cos(x) e^{2x} + 2 \sin(x) e^{2x} + C$$

Thus, we have

$$\int \sin(x) e^{2x} dx = \boxed{\frac{-\cos(x) e^{2x} + 2 \sin(x) e^{2x}}{5} + C}$$

4. Determine the following antiderivatives. Use the back of the previous page if you need more space.

15 pts

(a) $\int \sin^3(x) dx$

Solution: We use the identity $\sin^2(x) = 1 - \cos^2(x)$ to get

$$\int \sin^3(x) dx = \int (1 - \cos^2(x)) \sin(x) dx.$$

Now take $u = \cos(x)$ and $du = -\sin(x) dx$, giving

$$\int \sin^3(x) dx = -\int (1 - u^2) du = -u + \frac{u^3}{3} + C = \frac{\cos^3(x)}{3} - \cos(x) + C$$

15 pts

(b) $\int \frac{1}{\sec(4x)} dx$

Solution:

$$\int \frac{1}{\sec(4x)} dx = \int \cos(4x) dx = \frac{1}{4} \sin(4x) + C$$

15 pts


(c) $\int \frac{1}{x^2 \sqrt{x^2 - 1}} dx$

Solution: Take $x = \sec \theta$ so $dx = \sec \theta \tan \theta d\theta$. Then we have

$$\int \frac{1}{x^2 \sqrt{x^2 - 1}} dx = \int \frac{\sec \theta \tan \theta d\theta}{\sec^2 \theta \sqrt{\sec^2 \theta - 1}} = \int \frac{\tan \theta d\theta}{\sec \theta \sqrt{\tan^2 \theta}} = \int \frac{d\theta}{\sec \theta} = \int \cos \theta d\theta$$

This means we have $\sin \theta + C$ as our answer, but of course we need the answer in terms of x . Recall that we took $x = \sec \theta$, and so $\sin \theta = \frac{\sqrt{x^2 - 1}}{x}$ (see figure). Thus, we have shown

$$\int \frac{1}{x^2 \sqrt{x^2 - 1}} dx = \frac{\sqrt{x^2 - 1}}{x} + C$$



5. Evaluate these definite integrals. Use the back of the previous page if you need more space.

15 pts

(a) $\int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{1}{1-x^2} dx$

Solution: We use partial fractions:

$$\frac{1}{1-x^2} = \frac{A}{1+x} + \frac{B}{1-x}$$

so $1 = A(1-x) + B(1+x)$. Thus

$$A + B = 1 \quad -A + B = 0 \quad \text{hence} \quad A = \frac{1}{2}, B = \frac{1}{2}$$

$$\begin{aligned} \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{1}{1-x^2} dx &= \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{1/2}{1+x} + \frac{1/2}{1-x} dx = \frac{1}{2} \ln|1+x| - \frac{1}{2} \ln|1-x| \Big|_{-1/2}^{1/2} \\ &= \frac{1}{2} \left[\ln\left(\frac{3}{2}\right) - \ln\left(\frac{1}{2}\right) - \ln\left(\frac{1}{2}\right) + \ln\left(\frac{3}{2}\right) \right] \\ &= \ln\left(\frac{3}{2}\right) - \ln\left(\frac{1}{2}\right) = \boxed{\ln(3)} \end{aligned}$$

15 pts

(b) $\int_{-100}^{100} \frac{\sin^{21}(x)}{1+e^{x^2}} dx$

Solution: Since $\frac{\sin^{21}(x)}{1+e^{x^2}}$ is an odd function and the bounds are symmetric with respect to 0, the value of the integral is $\boxed{0}$.

15 pts

(c) $\int_0^1 \frac{x}{\sqrt{4-x^2}} dx$

Solution: Let $u = 4 - x^2$ so that $du = -2x dx$. When $x = 0$, $u = 4$ and when $x = 1$, $u = 3$. Thus we have

$$\int_0^1 \frac{x}{\sqrt{4-x^2}} dx = - \int_4^3 \frac{du}{2\sqrt{u}} = -\sqrt{u} \Big|_4^3 = -\sqrt{3} + \sqrt{4} = \boxed{2 - \sqrt{3}}.$$

6. Since $\int_0^1 \frac{1}{1+x^2} dx = \arctan(1) = \frac{\pi}{4}$, evaluating the integral $\int_0^1 \frac{4}{1+x^2} dx$ gives π .

20 pts

(a) Use Simpson's rule with 2 intervals to estimate $\int_0^1 \frac{4}{1+x^2} dx$.

Solution: Since there are two intervals, the width of each is $1/2$. Thus, Simpson's rule gives:

$$\frac{1}{3} \cdot \frac{1}{2} \left(f(0) + 4f(1/2) + f(1) \right) = \frac{1}{6} \left(4 + 4 \left(\frac{4}{1+1/4} \right) + 2 \right) = \boxed{\frac{94}{30}} \approx 3.13333$$

20 pts

(b) How many intervals are needed to estimate $\int_0^1 \frac{4}{1+x^2} dx = \pi$ within .0001 using the trapezoid rule?¹

Solution: We use the information in the footnote. We need to determine n so that

$$\frac{1}{12n^2} K \leq .0001$$

where K is the maximum of the absolute value of the second derivative of $4/(1+x^2)$ for x between 0 and 1. Since $\left| \frac{4(6x^2-2)}{(1+x^2)^3} \right|$ is a decreasing function on this interval, the maximum occurs at $x=0$, so we take $K = |-8/1| = 8$.

To solve $\frac{8}{12n^2} \leq .0001$, we multiply both sides by $10000n^2$ to get

$$\frac{80000}{12} \leq n^2,$$

so n is the smallest integer bigger than $\sqrt{20000/3} \approx 81.6$.

Thus, $\boxed{n = 82}$.

¹Use the following estimate for E_T using n intervals: If $|f''(x)| \leq K$ then $E_T \leq K \frac{(b-a)^3}{12n^2}$.

If $f(x) = \frac{1}{1+x^2}$, then $f''(x) = \frac{6x^2-2}{(1+x^2)^3}$